

## Drug Facts

### Active ingredients Purpose (in each tablet)

Aspirin 500 mg (NSAID\*).....Pain reliever  
Caffeine 60 mg .....Alertness aid/  
Pain reliever aid

\*Nonsteroidal anti-inflammatory drug

### Uses

- for the temporary relief of minor aches and pains associated with a hangover
- helps restore mental alertness or wakefulness when experiencing fatigue or drowsiness associated with a hangover
- also for the temporary relief of headaches or body aches and pains alone

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling
- asthma (wheezing) • shock

### Stomach bleeding warning:

This product contains a nonsteroidal anti-inflammatory drug (NSAID),

### Drug Facts (continued)

which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** • if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer • in children under 12 years of age

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- diabetes
- gout
- arthritis

### Drug Facts (continued)

#### When using this product

- the recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.
- for occasional use only. Do not use for more than 2 days for a hangover unless directed by a doctor. Not intended for use as a substitute for sleep. If fatigue or drowsiness persists or continues to occur, consult a doctor.

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain lasts for more than 10 days or gets worse
  - new symptoms occur
  - redness or swelling is present
  - ringing in the ears or a loss of hearing occurs

**If pregnant or breastfeeding** ask a health professional before use. It is especially

### Drug Facts (continued)

important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- upon waking, fully dissolve 2 tablets in 16 oz. of water and drink
- do not exceed recommended dosage

<b>Adults and children 12 years and over</b> (up to 60 years of age)	2 tablets every 6 hours, as needed, or as directed by a doctor. Do not exceed 8 tablets in 24 hours
--	---

<b>Adults 60 years and over</b>	2 tablets every 6 hours, as needed, or as directed by a doctor. Do not exceed 4 tablets in 24 hours
---------------------------------	---

<b>Children under 12 years</b>	Do not use
--------------------------------	------------

### Drug Facts (continued)

#### Other information

- each tablet contains: sodium 406 mg
- phenylketonurics: contains phenylalanine 12.6 mg per tablet
- store at room temperature (59–86°F). Protect from excessive heat.

#### Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

#### Questions or Comments?

1-800-970-1793 (weekdays)

**DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED**