

Common name	<i>Latin name</i>	Population	Isolation	Pollintation	Growth
<b>Umbelliferae</b>					
Parsnip	<i>Pastinaca sativa</i>	50-100	.5 miles	Outbreeder-insects	Biennial
Dill	<i>Anethum graveolens</i>	+/-50	.5 miles	Outbreeder-Insects	Annual
Carrot	<i>Daucus carota</i>	50-100	.5 miles	Outbreeder-insects	Biennial
Parsley	<i>Petroselinium crispum</i>	50-100	.5 mile	Outbreeder-insects	Biennial
Cilantro	<i>Coriandrum sativum</i>	50-100	.5 miles	Outbreeder-insects	Annual
<b>Brassicaceae</b>					
Cauliflower, Kale, Cabbage, Broccoli, Brussel sprouts	<i>Brassica oleraceae</i>	50-100	.5 miles	Outbreeder-insects	Biennial/Annual
Bok Choy, Turnips, Tatsoi Mustard, Mizuna	<i>Brassica rapa</i>	50-100	.5 miles	Outbreeder-insects	Annual
Giant Red Mustard, Green Wave Mustard	<i>Brassica juncea</i>	50-100	.5 miles	Outbreeder-insects	Annual
Arugula	<i>Eruca sativa</i>	50-100	.5 miles	Outbreeder-insects	Annual
Radish	<i>Raphanus sativus</i>	50-100	.5 miles	Outbreeder-insects	Annual
Garden Cress	<i>Lepidium sativum</i>	50-100	.5 miles	Outbreeder-insects	Annual
<b>Chenopodaceae</b>					
Beet	<i>Beta vulgaris</i>	50-100	2 miles	Outbreeder-wind	Biennial
Chard	<i>Beta vulgaris</i>	50-100	2 miles	Outbreeder-wind	Biennial
Spinach	<i>Spinacia oleracea</i>	50-100	1 miles	Outbreeder-wind	Annual dioecious
Orach	<i>Atriplex hortensis</i>	50-100	.k5 miles	Outbreeder-wind	Annual
<b>Compositae</b>					
Lettuce	<i>Lactuca sativa</i>	+/-20	10ft	Inbreeder	Annual
<b>Leguminosae</b>					
Pea	<i>Pisum sativum</i>	+/-20	10ft	Inbreeder	Annual