

## RADON AND HEALTH

### *What is radon and where does it come from?*

Radon is a natural radioactive gas without odour, colour or taste. It cannot be detected without special equipment. Radon occurs as a product of uranium decay. Uranium is a natural radioactive material found in varying amounts in all rocks, soil, concrete and bricks. It occurs everywhere on earth, especially in rocky and mountainous areas. Radon is an unstable radionuclide that disintegrates through short lived decay products before eventually reaching the end product of stable lead. The short lived decay products of radon are responsible for most of the hazard by inhalation.

Radon and its decay products called radon daughters or radon progeny emit highly ionizing alpha-radiation. Decay products are suspended in the air which we breathe. Although the risk is very low when radon is diluted to extremely low concentrations in the open, radon in room air typically contributes up to 50% to the background radiation (fig 1.). However, in places such as caves and mines, it can accumulate up to dangerous concentrations and may cause substantial health damage after long-term exposure. Radon can also be found in drinking water and this can sometimes present a hazard. Certain types of geology, such as granite and volcanic soils, as well as aluminous shales, are more likely to contain radon. Conversely, low concentrations of this gas are expected in sedimentary rocks.




Figure 1. Sources and average distribution of natural background radiation for the world population (UNSCEAR, 2000).

### *How radon is measured*

Radioactivity of a radionuclide, for instance radon, is reported in becquerels, Bq.

**1 becquerel (1 Bq) = 1 disintegration of atom per second**

Radon concentrations in the air are measured as the amount of radioactivity (Bq) in a cubic metre of air ( $\text{Bq}/\text{m}^3$ ). The average radon level outdoors is  $5 \text{ Bq}/\text{m}^3$  and in the homes of the United Kingdom, for instance, is  $20 \text{ Bq}/\text{m}^3$  of air.

### *How does radon enter houses?*

Radon gas enters houses from the ground through cracks in concrete floors and walls, through gaps between floor and slab, and around drains and pipes, and small pores of hollow-block walls (fig.2). Consequently, radon levels are usually higher in basements, cellars and ground floors. Depending on a number of factors, the concentration of radon indoors varies with the time of the year, from day to day, and from hour to hour. Because of this time-variation, reliable measurements of mean concentrations in air should be made for at least three months. The level of radon in homes can be measured with simple device that are returned to the testing company to interpret the results.




Figure 2. Radon entry ways

### *Health effects of radon*

The adverse health effects of exposure to radon are caused primarily by damage due to alpha-particles. The possible effects will depend on exposure level. The main danger from high radon exposure is an increased risk of lung cancer. Radon as a noble gas is rapidly exhaled after being breathed in; however, radon progeny, combine with other molecules in the air and with particles of dust, aerosols or smoke, and readily deposit in the airways of the lung. While lodged there, the progeny emit ionizing radiation in the form of alpha particles, which can damage the cells lining the airways. Experiments have confirmed that ionizing radiation affecting bronchial epithelial cells could cause cancer.

Epidemiological studies on thousands of uranium miners in different countries, including Germany, USA, Canada, Czechoslovakia, and others, also support this fact. Analyses

have been undertaken of several miner studies, totalling 68,000 men of whom 2,700 died from lung cancer. An increased risk of all histological types of lung cancer, including small cell carcinoma, adenocarcinoma, and squamous cell carcinoma, has been associated with occupational exposure to radon. But exposure to radon in houses can also lead to lung cancer. It is believed, for example, that every year more than 15 000 deaths from lung cancer occur due to radon exposure in the United States and more than 2 500 deaths in the United Kingdom.



**Lung cancer risk is much higher when radon exposure is combined with smoking. According to the BEIR IV report of the US National Academies of Sciences, for men exposed to radon at work, smokers were 10 times more likely to get lung cancer risk than non-smokers.**

The risk of lung cancer from radon in home can be estimated by two ways: by direct epidemiological studies of residential radon and by projecting occupational risk estimates to lower levels of radon in homes. The uncertainties associated with residential studies are larger than those with miner studies, primarily because the risk is smaller at the low exposures encountered in most homes. It is also difficult to estimate the radon exposures that people have received over their lifetimes, probably in several houses. Despite the uncertainties, it is clear that far more lung cancers are caused by smoking than are caused by radon.

It is believed that the relationship between radon and risk of lung cancer is linear. In other words, doubling the exposure doubles the risk and halving the exposure halves the risk. Doubling of the risk means much more for a smoker, who is already at high risk of lung cancer, than for a non-smoker with a very small base line risk. Lung cancer risk from residential radon exposure is substantially lower since the exposure in homes is much lower than in mines, although the risk increases with radon concentration level and duration of exposure. For life-time exposure to radon of 20 Bq/m<sup>3</sup> level at home the risk of lung cancer is estimated to be 0.3% (or 3 deaths in 1000 people). For comparison, risk of accidental death at home is 0.7% (or 7 in 1000).

It has been suggested that other effects of radon exposure include increased risk of non-malignant respiratory diseases but this is much less clearly established than the lung cancer risk. It is still not clear whether children are more sensitive to radon exposure. Studies on childhood leukaemia (the most common form of cancer in childhood) have not found clear evidence of risk associated with radon concentrations in homes.

### ***Radon in drinking water***



Ninety-five percents (95%) of exposure to radon is from indoor air; about one percent (1%) is from drinking water sources. Most of this 1% drinking water exposure is from inhalation of radon gas released from running water activities, such as bathing, showering, and cleaning. Only 0.1% of exposure is from swallowing drinking water contaminated with radon gas. Although the effects of ingested radon are not fully understood, calculations suggest that the great majority of the radiation dose from such exposure is in the stomach.

In many countries, some homes obtain drinking water from groundwater sources (springs, wells, adits, and boreholes). Because of its volatility, radon contamination occurs primarily in ground, not surface, water sources of drinking water. Underground, this water often moves through rock containing natural uranium that releases radon to the water. That is why, water from wells normally has much higher concentrations of radon than surface water such as rivers, lakes, and streams. The U.S. National Academy of Sciences reports that the average concentration of radon in public water supplies derived from surface waters is usually less than 0.4 Bq/l, and it is about 20 Bq/l in ground water sources in the United States.

### *Radon in the air is easy to detect*

It is possible for one house to have elevated levels of radon while a neighbouring one does not. Measurement is the only reliable way to determine levels of radon in a house. Measurements are normally carried out using passive detectors which are left in the home for periods from days to months. Because radon levels vary from day to day and from season to season, measurements over several months are better than short-term measurements for estimating annual average radon levels.

Detectors can be purchased from testing companies, but must be returned to them for assessment. To accurately reflect people's true exposure to radon, detectors must be placed in rooms where people spend most of the time (living room, bedrooms, offices). Instructions should be followed carefully when installing the detectors. They should not be placed in rooms that are rarely used.

There are three main types of passive radon detector:

- Etched track (or alpha track) detectors are normally placed for a period of one to 12 months.
- Electret ion chambers are available with different sensitivities, some being suitable for measurements over a few days and some being suitable for measurements over months.
- Charcoal detectors measure average radon levels over a few days, so give less accurate estimates of annual average radon levels. They may be used for screening measurements.

A list of suppliers of radon detectors may be obtained from the national or regional health authorities or from research workers or specialists in this field.

### *Action level of radon concentration in the air and water*

Many countries have defined an Action Level of radon concentration to guide their program to control domestic exposure to radon. The Action Level is not a boundary between safe and unsafe, but rather a level at which action on reduction of radon level will usually be justified. Some people may choose to take action when the Action Level is approached. For example, many countries consider radon concentration in the air of 200 Bq/ m<sup>3</sup> as an Action Level at which mitigation measures should be taken to reduce radon level in homes. Guideline values (action levels) of radon vary among countries (Table 1).

The United States set a Maximum Contaminant Level for radon in drinking water of 150 Bq/l for radon concentration in drinking water from private water supplies, The European Union Commission recommended Action Level of 1000 Bq/l. This is set so that the risk to a typical person drinking such water is similar to the risk from breathing air which contains radon at the Action Level of 200 Bq/m<sup>3</sup>.

Table 1. Domestic radon concentrations and Action Levels in different countries

Country	Average radon concentration in homes (Bq/m <sup>3</sup> )	Action Level (Bq/m <sup>3</sup> )
Czech Republic	140	200
Finland	123	400
Germany	50	250
Ireland	60	200
Israel	*	200
Lithuania	37	100
Luxembourg	*	250
Norway	51-60	200
Poland	*	400
Russia	19-250	*
Sweden	108	400
Switzerland	75	400
United Kingdom	20	200
European Community	*	400
USA	46	150
Canada	*	800

\* not available at the moment

### ***Radon may be reduced at Your Home***

Radon levels in room air can be lowered in a number of ways, from sealing cracks in floors and walls to changing the flow of air into the building. The five principle ways of reducing the amount of radon entering a house are:

- Improving the ventilation of the house;
- Sealing floors and walls;
- Increasing under-floor ventilation;
- Installing a radon sump system;
- Installing a whole house positive pressurization or positive supply ventilation system.

Some of these solutions are not suitable for all types of houses, nor are they suitable for all levels of radon. In some cases, more than one solution is needed in resolving the radon problem.

Radon safety needs to be considered when new houses are built. In many countries, for instance in England and in the United States, including protective measures in new buildings has become a routine procedure. These measures have been proven to keep radon levels well below the recommended action level, both in the short and long term.

### *Related Information Links on the Web*

More information on Radiation, Radon, Health Effects of Radon, Radon Measurements and Mitigation could be found on the following websites:

United Kingdom National Radiation protection Board

<http://www.nrpb.org>

US Environmental Protection Agency

<http://www.epa.gov/iaq/radon/index.html>

Swiss Public Health Department

<http://www.ch-radon.ch>

US National Radon Safety Board

<http://www.nrsb.org/>

US National Research Council, Board on Effects of Ionising Radiation VI Report on Health Effects of Exposure to Radon (full text)

<http://www.nap.edu/books/0309056454/html/index.html>

Find state Radon Service Officials in the U.S.A.

<http://www.epa.gov/iaq/contacts.html>

US Nuclear Regulatory Committee

<http://www.nrc.gov>

American Lung Cancer Association

[http://www.lungusa.org/air/radon\\_factsheet99.html](http://www.lungusa.org/air/radon_factsheet99.html)

Environmental Radon Newsletter published by NRPB, UK

[http://www.nrpb.org/publications/newsletters/environmental\\_radon/index.htm](http://www.nrpb.org/publications/newsletters/environmental_radon/index.htm)

Building Research Establishment, Foundation for the Built Environment, UK

[www.bre.co.uk](http://www.bre.co.uk)

Health Canada: Radon

<http://www.hc-sc.gc.ca/ehp/ehd/catalogue/general/iyh/radon.htm>