

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 09/2018

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:ADONA TABLETS 30mg

Active ingredient:Carbazochrome sodium sulfonate hydrate

Dosage form:orange-yellow to orange-yellowish brown tablet, diameter: 8.0 mm, thickness: 3.6 mm

Imprint or print on wrapping: (Face) Adona30mg, アドナ 30mg, アドナ 30, TA107
(Back) アドナ 30mg



Effects of this medicine

This medicine shortens bleeding time and shows hemostatic action by acting to blood capillary to inhibit increased blood vessel permeability and reinforce vessels resistance.

It is usually used to treat bleeding tendency (purpura, etc.) caused by attenuation of microvascular resistance and increased capillary permeability, bleeding from skin, mucosa or inner membrane, fundal/renal/uterine bleeding and intraoperative or postoperative abnormal bleeding.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 10 to 30 mg of the active ingredient at a time, 3 times daily. The dosage should be adjusted according to your age and symptoms. Strictly follow the instructions.
- If you miss a dose, take a dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- The urine may change color to orange-yellow.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include , , and . If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.