

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 11/2017

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:ALESION Ophthalmic Solution 0.05%

Active ingredient:Epinastine hydrochloride

Dosage form:limpid and colorless, ophthalmic solution, 5mL, (bottle) clear and colorless, (cap) deep yellow

Print on wrapping:



Effects of this medicine

This medicine suppresses allergic symptoms such as itchy/bloodshot eyes by histamine H₁-receptor antagonism and suppressing release of allergy-causing substances such as histamine.

It is usually used to treat allergic conjunctivitis.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines. If you wear contact lenses.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply 1 drop at a time, 4 times a day (morning, afternoon, evening, and before bedtime). Strictly follow the instructions.
- Before applying, wash your hands thoroughly with soap and running water. Pull down your lower eyelid so that the dropper tip does not touch eye or eyelashes and apply the medicine. After applying, do not blink and keep your eyes closed for a while. Wipe off any excess liquid with a clean gauze or tissue immediately. If you need to apply other eye drops at the same time, wait more than 5 minutes.
- If you miss a dose, apply the missed dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include stinging eyes, foreign body sensation in eyes, photophobia, blepharitis (erosion of eyelids), and eye mucus. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of reach of children. Store away from direct sunlight and heat.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.