

Drug Information Sheet("Kusuri-no-Shiori")

Internal
Revised: 10/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: ALLELOCK OD Tablets 2.5**Active ingredient:** Olopatadine hydrochloride**Dosage form:** very light yellow tablet, diameter: 7.5 mm, thickness: 2.9 mm**Imprint or print on wrapping:** KH 022 2.5mg アレロック OD**Effects of this medicine**

This medicine suppresses production and release of allergic substances (histamine, etc.) in the body. It also resolves and relieves allergic symptoms in skin and nose by inhibiting activities of these allergic substances. It is usually used in the treatment of allergic rhinitis, hives, and itch associated with skin diseases.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
If you have renal disorder or hepatic disorder.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 2 tablets (5 mg of the active ingredient) at a time, twice daily, in the morning and before bedtime. The dosage may be adjusted according to the disease, age or symptoms. For children 7 years old or over, take 2 tablets (5 mg) at a time, twice daily, in the morning and before bedtime. Strictly follow the instructions.
- You may swallow the medicine either with water as an ordinary tablet or after dissolving with saliva in the mouth.
- If you miss a morning dose and notice it until noon, take a dose as soon as possible. However, if you notice it late in the day, skip the missed dose and continue your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- The medicine may cause sleepiness. Avoid driving a car or operating dangerous machinery.
- If you take the medicine for treatment of pollinosis, you need to take the medicine until the pollen season ends.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include sleepiness, malaise, dry mouth, abdominal discomfort, headache, dull headache and dizziness. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- general malaise, loss of appetite, nausea, yellowing in skin and white of eyes [fulminant hepatitis, hepatic dysfunction, jaundice]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.