Drug Information Sheet("Kusuri-no-Shiori")

External evised: 01/2021

	Revised: 01/2021
The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response,	
patients should understand their medication and cooperate with the treatment.	
Brand name: ANTEBATE LOTION 0.05%	
Active ingredient: Betamethasone butyrate propionate	
Dosage form:white lotion	
Imprint or print on wrapping:アンテベートローション 0.05%, TO-186L	ALC: NO
Effects of this medicine	. 1
This medicine is a steroid for external use, and suppresses skin inflammation and improves	symptoms such as
redness, swelling and itch.	
It is usually used to treat eczema, dermatitis, and psoriasis.	
The following patients may need to be careful when using this medicine.Be sure t	o tell your doctor and
pharmacist.	
• If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicine	
If you have eczematous external otitis with eardrum perforation, skin infections due to bacteria/fungi, ulcer and	
burn/frostbite.	
• If you are pregnant or breastfeeding.	
• If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal	
effects. Beware of over-the-counter medicines and dietary supplements as well as other	prescription medicines.)
Dosing schedule (How to take this medicine)	1.1 0 1)
• Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))	
•In general, apply an appropriate amount of the medicine to the affected area, once to several times a day. Strictly	
follow the instructions.	
•Pay extra attention when you apply this medicine to areas that touch the diaper. Follow the instructions.	
•Do not apply this medicine to your eyes (cornea and conjunctiva). If the medicine accidentally gets into your eyes,	
rinse it off with water. (If you still feel something wrong, consult with your doctor or pharmacist.)	
• If you miss a dose, apply the missed dose when you remember. However, if it is almost time for the next dose, skip	
the missed dose and then apply the next dose as instructed. You should not apply two doses at one time.	
• If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.	
•Do not stop applying this medicine unless your doctor instructs you to do so.	
Precautions while taking this medicine	
• If you use this medicine for a long period of time, consult with your doctor regularly.	
•Do not use this medicine as a makeup base or after shaving.	
Possible adverse reactions to this medicine	
The most commonly reported adverse reactions include skin fungal infection (candidiasis, ti	
infection (contagious impetigo, folliculitis, etc.), acne, steroid skin (skin atrophy, small visil	
flushing, etc.), stimulating feeling and eczema. If any of these symptoms occurs, consult wit	h your doctor or
pharmacist.	
The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated	
in brackets. If any of these symptoms occur, stop taking this medicine and see yo	our doctor
immediately.	
•(When you use this medicine in a large dose or over a long period of time over a large area	
the eyelids) eye pain, sensitive to light, foggy vision, headache [ocular hypertension, glat	
The above symptoms do not describe all the adverse reactions to this medicine. Consult with your	
doctor or pharmacist if you notice any symptoms of concern other than those listed above. Storage conditions and other information	
	poraturo (1 to 30 dograco
•Keep out of the reach of children. Store away from light and moisture. Store at room temperature (1 to 30 degrees Celsius) in this original bottle.	
•Discard the remainder. Do not store them. Consult with your pharmacy or medical institution on how to discard them.	
• Storing upside down is supposed to be easy to put out the lotion.	
For healthcare professional use only / /	

For further information, talk to your doctor or pharmacist.