

# Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 03/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:** APLACE Tablets 100mg

**Active ingredient:** Troxipide

**Dosage form:** white tablet, diameter: 8.1mm, thickness: 4.1mm

**Print on wrapping:** (face) アプレース, 錠 100mg, KP-296  
(back) APLACE, Tab. 100mg, アプレース, 100mg



**Effects of this medicine**

This medicine acts on gastric mucosa and helps repair tissues of gastric mucosa. It is usually used to treat gastritis/gastric ulcer.

**Before using this medicine, be sure to tell your doctor and pharmacist**

- If you ever experienced any allergic reaction (itch, rash etc.) to any medicine.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

**Dosing schedule (How to take this medicine)**

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, for adults, take 1 tablet (100mg of the active ingredient) at a time, 3 times a day after meals. The dosage may be adjusted according to your age and symptoms. Strictly follow the instructions.
- If you missed a dose, take a dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. DO NOT take a double dose to make up for the missed dose.
- If you took too much of this medicine (more than prescribed), consult with your doctor/pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

**Precautions while taking this medicine**

**Possible adverse reactions to this medicine**

The most commonly reported adverse reactions include constipation. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

- fever, feeling unwell, redness [shock, anaphylactoid symptoms]
- malaise, loss of appetite, yellowing of the skin/eyes [liver dysfunction, jaundice]

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

**Storage conditions and other information**

- Keep out the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.