

Drug Information Sheet("Kusuri-no-Shiori")

Internal
Revised: 05/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Acofide Tablets 100mg**Active ingredient:** Acotiamide hydrochloride hydrate**Dosage form:** white tablet, diameter 9.1 mm, thickness 4.2 mm**Imprint or print on wrapping:** (Face) アコファイド, 100mg (Back) Acofide100mg, アコファイド**Effects of this medicine**

This medicine improves gastrointestinal motility and promotes excretion of stomach contents by inhibiting acetylcholinesterase activities.

It is usually used for the treatment of postprandial fullness, upper abdominal bloating, and early satiation due to functional dyspepsia.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 1 tablet (100mg of the active ingredient) at a time, 3 times a day before meals. Strictly follow the instructions.
- If you miss a dose, skip the missed dose and follow your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine**Possible adverse reactions to this medicine**

The most commonly reported adverse reactions include diarrhoea, constipation, nausea, vomiting, abdominal pain, rash and hives. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.