

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 09/2023

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Astat Cream 1%

Active ingredient:Lanoconazole

Dosage form:white cream

Imprint or print on wrapping:アスタットクリーム 1% 10g



Effects of this medicine

This medicine shows antifungal effects by acting on fungal cell membrane.

It is usually used to treat athlete's foot/ringworm (tinea), candidiasis and tinea versicolor.

The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply to the affected area once a day. Strictly follow the instructions.
- Do not apply the medicine to severely eroded surfaces.
- Do not apply it to your eyes. If this medicine accidentally gets into your eyes, wash out your eyes with water immediately and consult with your doctor or pharmacist.
- If you miss a dose, apply as soon as possible. Do not apply a double dose to make up for the missed dose.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include skin inflammation, irritating sensation and redness. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store at room temperature (1 to 30 degrees Celsius), away from direct sunlight and moisture.
- Discard any unused leftover amount. Do not keep it. If you do not know how to dispose of them, please consult the pharmacy or medical institution where you received them. Do not give them to others.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.