

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 07/2022

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:CORECTIM Ointment 0.5%
Active ingredient:Delgocitinib
Dosage form:white ointment
Imprint or print on wrapping:コレクチム軟膏 0.5%



Effects of this medicine

This medicine inhibits enzymes called as Janus kinase to suppress the actions of factors involved in immuno/inflammatory actions such as lymphocyte activity and thus to suppress inflammation. It is usually used to treat atopic dermatitis.

The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, apply an appropriate amount of this medicine to the affected area twice a day. A single dose should not exceed 5 g at a time. In general, for children, apply an appropriate amount of 0.25% formulation to the affected area twice a day. Depending on symptoms, apply an appropriate amount of this medicine (0.5% formulation) to the affected area twice a day. The applied dose should not exceed 5 g at a time. For children, the dose may be adjusted according to the physique. In any case, strictly follow the instructions.
- If the symptoms do not improve after 4 weeks of treatment with this medicine, consult with your doctor.
- For children, this medicine may be replaced with a 0.25% formulation when the symptoms improve. Follow the instructions of your doctor or pharmacist.
- Apply by avoiding skin infected area.
- Do not apply to mucosa, injured skin or eroded area.
- If this medicine gets into your eyes by any chance, wash out with water immediately.
- If you miss a dose, apply the missed dose when you remember. You should not apply two doses at one time.
- If you accidentally use more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop using this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include folliculitis (eruption) and acne (pimple). If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them. If you do not know how to discard the remainder, ask your pharmacy or medical institution. Do not give the unused medicines to others.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.