

# Drug Information Sheet("Kusuri-no-Shiori")

Internal

Published: 07/2008


The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**Cephadol Tablets 25mg

**Active ingredient:**Difenidol hydrochloride

**Dosage form:**pale yellow tablets,  $\phi$  : 7.1mm, thickness: 3.2mm

**Print on wrapping:** (face) 108 (back) セファドール 25mg



## Effects of this medicine

This medicine is effective for improvement of circulation of vertebral artery, for control of vertebral nerve pathway, and for suppression of nystagmus.

Used dizziness caused by inner ear disturbance.

## Before using this medicine, be sure to tell your doctor and pharmacist

- If you ever experienced any allergic reaction (itch, rash etc.) to any medicine.
- If you have renal dysfunction.
- If you have glaucoma.
- If you have a history of drug eruption, urticaria, and so on.
- If you have obstructive disease in urinary tract such as prostatic hyperplasia.
- If you have obstruction in digestive tract.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is<< to be written by a healthcare professional>>
- General dosage regimen: Take 1 to 2 tablets (25 to 50mg of the active ingredient) at a time, 3 times daily, however, the dosage may be adjusted according to the age or symptoms. Strictly follow the instructions of your doctor/pharmacist.
- Take a dose as soon as possible when you remember that you missed a dose. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. DO NOT take a double dose to make up for the missed dose.
- If you took too much of this medicine (more than ordered), check with your doctor/pharmacist.
- Do not stop taking this medicine without the instructions of your doctor.

## Precautions while taking this medicine

## Possible adverse reactions to this medicine

Common side effects are reported as below. If any of them occurs, check with your doctor/pharmacist: Dry mouth, anorexia, heartburn, floating feeling, instability feeling, feeling of facial warmth, palpitation, hallucination, rash, urticaria, focus disorder of the eye, mydriasis, etc.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

No pertinent entries.

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

## Storage conditions and other information

- Keep out the reach of children. Store away from heat, moisture and direct sunlight.
- Do not keep the remains. Discard them.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.