

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Published: 07/2008

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Coldrin Tablets 12.5mg

Active ingredient:Clofedanol hydrochloride

Dosage form:white tablet, φ : 7.6 mm, thickness: 4.6 mm

Print on wrapping:(front) 218 (back) コルドリン錠 12.5mg



Effects of this medicine

This medicine acts on the cough center of the brain stem and shows antitussive action to suppress cough. It is usually used in the treatment of cough in acute bronchitis and acute upper respiratory tract inflammation.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you ever experienced any allergic reaction (itch, rash etc.) to any medicine.
If you are a patient with debility.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is<< to be written by a healthcare professional>>
- General dosage regimen: For adults, take 2 tablets (25 mg of the active ingredient) at a time, three times daily. The dosage may be adjusted according to the age or symptoms. Strictly follow the instructions of your doctor/pharmacist.
- Take a dose as soon as possible. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. DO NOT take double doses to make up for the missed dose.
- If you took too much of the medicine (more than ordered), check with your doctor/pharmacist.
- Do not stop taking the medicine without the instructions of your doctor.

Precautions while taking this medicine

Possible adverse reactions to this medicine

Common side effects are reported as below. If any of them occurs, check with your doctor/pharmacist: nausea, vomiting, loss of appetite, dizziness, rash, itch, etc.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- hives, cold sweat, breathing difficulty [shock, anaphylactoid symptoms]
- severe stomatitis, red rash with swelling in center, congested eyelids, bloodshot eyes [oculomucocutaneous syndrome]
- fever, joint pain, rash and blisters with pain [erythema multiforme exudativum]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.