

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 10/2018

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Dalacin T Lotion 1%

Active ingredient: Clindamycin phosphate

Dosage form: limp and colorless lotion

Print on wrapping: ダラシン T ローション 1%



Effects of this medicine

This medicine is a topical lincomycin antibiotic and shows bactericidal effect by inhibiting protein synthesis of bacteria.

It is usually used to treat acne (with suppurative inflammation).

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply appropriate amount to the affected area after washing your face, twice a day. Strictly follow the instructions.
- The medicine should be used only for dermal application. Not for ophthalmic use.
- If you miss a dose, skip the missed dose and continue your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include irritating sensation and erythema. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- abdominal pain, frequent diarrhea with mucus/bloody stool, fever [serious colitis with bloody stool such as pseudomembranous colitis]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only

/ /

For further information, talk to your doctor or pharmacist.