

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 11/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:ECLAR OINTMENT 0.3%

Active ingredient:Deprodone propionate

Dosage form:white to faint yellow ointment

Imprint or print on wrapping:HP2100-O



Effects of this medicine

This medicine is a steroid preparation with anti-inflammatory properties and suppresses symptoms such as itching and swelling.

It is usually used to treat eczema/dermatitis, drug rash, insect bites, prurigo, psoriasis, erythroderma, erythema, palmoplantar pustulosis, purpura and alopecia areata.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itching, rash, etc.) to any medicines or foods. If you have skin infection due to bacteria or fungi, skin diseases caused by insects (scabies, hair lice, etc.), eczematous external otitis with eardrum perforation, ulcer or burn/frostbite.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply an adequate amount of the medicine to the affected area, once to several times a day. Strictly follow the instructions.
- Do not apply the medicine to your eyes.
- For infants and children, avoid applying for a long period/mass and avoid applying in sealed states (under a tight-fitting diaper, etc.).
- If you miss a dose, apply the missed dose when you remember. If it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- Do not apply this medicine under makeup or after shaving.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include irritating sensation of skin, itching, contact dermatitis, folliculitis and rash. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- severe eye pain, foggy vision, rapid reduction of visual acuity [glaucoma, posterior capsule cataract]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from light, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.