Drug Information Sheet("Kusuri-no-Shiori")

External
Revised: 01/2018

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Eurax Cream 10%

Active ingredient: Crotamiton
Dosage form: white to yellowish white cream
Print on wrapping: CG EX (written on tube)

Effects of this medicine

This medicine gives mild burning sensation to the skin, and consequently reduces itching of skin due to eczema or hives.
It is usually used to treat eczema, hives, dermal pruritus, etc.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply or rub a proper amount of the medicine on the affected site several times a day. Strictly follow the instructions.
  This medicine is designed for topical application to the skin. Never take it orally.
  Do not apply on mucous membranes such as lips, in or around the eyes. If the medicine got in your eyes, wash out with water or lukewarm water immediately.
  If the medicine is used for children, use it under supervision of the parent. Do not apply it to a wide skin area.
  You may feel warmth sensation just after applying the medicine, which will disappear in a short time.
- If you miss a dose, apply a dose as soon as possible. You should not use two doses at one time.
- If you accidentally use more than your prescribed dose, wipe off any excess of the medicine and consult with your doctor or pharmacist.
- Do not stop using this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include feeling of skin irritation (feeling hot, irritated sensation), contact dermatitis (redness, etc.), itch, rash, eczema, erythema and angioedema. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Never transfer the medicine to other container to keep its quality.
- Keep out the reach of children. Store away from heat.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.