

Drug Information Sheet("Kusuri-no-Shiori")

Internal
Revised: 07/2011

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Eviprostat Tablets DB

Active ingredient: Herbal extracts mixture

Dosage form: white tablet, diameter 7.8 mm, thickness 5.0 mm

Imprint or print on wrapping: (face) 222, (back) エビプロスタット配合錠 DB



Effects of this medicine

This medicine has anti-inflammatory effect and diuretic effect. It also improves enlargement of the prostatic tissue and sterilizes the urinary tract.

It is used to treat dysuria, residual urine/feeling of residual urine, and pollakiuria associated with prostatic hyperplasia.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you ever experienced any allergic reaction (itch, rash etc.) to any medicines or foods.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- General dosage regimen: For adults, take 1 tablet, 3 times a day. However, the dosage may be adjusted according to the symptoms. Strictly follow the instructions of your doctor/pharmacist.
- Take the missed dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. DO NOT take double doses to make up for the missed dose.
- If you took too much of the medicine (more than ordered), check with your doctor/pharmacist.
- Do not stop taking the medicine without the instructions of your doctor.

Precautions while taking this medicine

Possible adverse reactions to this medicine

Common side effects are reported as below. If any of them occurs, check with your doctor/pharmacist: loss of appetite, abdominal pain, gastric discomfort, hypersensitivity symptoms (such as rash, itching), erythema multiform (circular or oval eruption), jaundice, etc.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, high temperature and humidity.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.