

# Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 12/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**HEPARINOID LOTION 0.3% "Nichiiiko"

**Active ingredient:**Heparinoid

**Dosage form:**colorless to faint yellow limpid lotion

**Imprint or print on wrapping:**へパリン類似物質ローション 0.3%「日医工」



## Effects of this medicine

This medicine keeps moisture of the skin to relieve symptoms of dry skin. It also accelerates blood flow of the applied area to relieve pain or swelling caused by vascular insufficiency.

It is usually used to treat and prevent asteatosis, keratoderma tylodes palmaris progressiva, chilblain, hypertrophic scar/keloid, and to treat pain and inflammatory disease caused by vascular insufficiency (induration and pain after injection), thrombophlebitis (including hemorrhoids), posttraumatic (bruise/sprain/contusion) swelling/hematoma/tenosynovitis/muscle pain/arthritis and muscular torticollis (infant).

## The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.  
If you have a hemorrhagic blood disease (hemophilia, thrombocytopenia, purpura, etc.).  
If you are expected to have a serious result with slight bleeding.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, apply an appropriate amount of lotion to the affected area once to several times a day. Strictly follow the instructions.
- Avoid applying this medicine directly on ulcer or erosion.
- Take care not to let the medicine get in your eyes.
- If you miss a dose, apply a dose as soon as possible. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

## Precautions while taking this medicine

### Possible adverse reactions to this medicine

The most commonly reported adverse reactions include skin irritation, dermatitis, itch, redness, rash, flush and purpura. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

No pertinent entries.

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

### Storage conditions and other information

- Close the cap and store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.