

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 07/2023

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Hirudoid Cream 0.3%

Active ingredient:Heparinoid

Dosage form:white cream

Imprint or print on wrapping:



Effects of this medicine

This medicine relieves dry skin due to its moisturizing effect. It also treats pain or swelling by promoting blood circulation.

It is usually used to treat conditions such as dry skin (seborrheic dermatitis), chapped hands, chilblains, keloids, or hypertrophic scars, etc.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you ever experienced any allergic reaction (itch, rash etc.) to any medicine or foods.
If you have the following disease: hemophilia, thrombocytopenia, purpura, etc.
If even slight bleeding can lead to serious consequences (this medicine might make it harder for the blood to clot).
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, rub an appropriate amount of this medicine into the affected area, or apply the gauze previously spread with the medicine to the affected area once to several times a day. Strictly follow the instructions of your doctor/pharmacist.
- Do not apply this medicine to areas where the skin is sored or wounded.
- If you miss a dose, apply a single dose as soon as you remember. Do not apply a double dose to make up for the missed dose.
- If you applied too much of this medicine (more than ordered), check with your doctor/pharmacist.
- Do not stop applying this medicine without the instructions of your doctor.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include dermatitis, itch, redness, rash, hot flush, skin irritation and purpura. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them. If you do not know how to dispose of them, please consult the pharmacy or medical institution where you received them. Do not give the unused medicines to others.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.