

# Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 09/2017


The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**Hirudoid Soft Ointment 0.3%

**Active ingredient:**Heparinoid

**Dosage form:**white cream

**Print on wrapping:**



## Effects of this medicine

This medicine keeps moisture of the skin, to relieve symptoms of dry skin. It also accelerates blood flow of the applied area to lessen pain or swelling caused by blood circulation disorder.

Usually, this medicine is used in the treatment of chilblain, hypertrophic scar, keloid, keratoderma tylodes palmaris progressiva or asteatosis etc.

## Before using this medicine, be sure to tell your doctor and pharmacist

- If you ever experienced any allergic reaction (itch, rash etc.) to any medicine. If you have a hemorrhagic blood disease (hemophilia, thrombocytopenia, purpura). If you have a serious result even with a slight bleeding.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, rub an appropriate amount of this medicine into the affected area, or apply the gauze previously spread with the medicine to the affected area once to several times a day. Strictly follow the instructions of your doctor/pharmacist.
- Do not apply this medicine to your eyes or open wounds (ulcer, erosion).
- If you miss a dose, apply as soon as possible. Do not apply a double dose to make up for the missed dose.
- If you applied too much of this medicine (more than prescribed), check with your doctor/pharmacist.
- Do not stop applying this medicine without the instructions of your doctor.
- Do not apply a double dose to make up for the missed dose, apply as soon as possible.
- If you applied too much of this medicine (more than ordered), check with your doctor/pharmacist.
- Do not stop applying this medicine without the instructions of your doctor.

## Precautions while taking this medicine

### Possible adverse reactions to this medicine

The most commonly reported adverse reactions include dermatitis, itch, redness, rash, hot flush, irritation and purpura. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

No pertinent entries.

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

### Storage conditions and other information

- Keep out of the reach of children. Store away from heat, moisture and direct sunlight.
- Discard any unused leftover amount. Do not keep it.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.