

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 07/2015

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Hyaluronate PF Ophthalmic Solution 0.1% "NITTEN"

Active ingredient: Purified sodium hyaluronate

Dosage form: limpid and colorless, viscous, eye solution, (cap) very light blue, (bottle) white translucent, (accompanying bag) translucent



Print on wrapping:

Effects of this medicine

This medicine promotes corneal injury healing by promoting adhering/repairing corneal epithelial cells and prevents eyes from dry out by retaining/stabilizing tears.

It is usually used to treat endogenous keratoconjunctival epithelial disorder such as Sjogren's syndrome, Stevens-Johnson syndrome and dry eye, and exogenous keratoconjunctival epithelial disorder (eye scratch etc.) associated with after surgery, drug-induced, injury and wearing contact lenses.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
If you wear contact lenses.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply 1 drop of 0.1% preparation at a time, 5 to 6 times a day. The dosage may be adjusted according to your symptoms. If there is not enough effect, apply 0.3% preparation in the same way. This medicine is 0.1% preparation. Strictly follow the instructions.
- Before using, wash your hands with soap and water. Tilt your head back, pull down your lower eyelid, open your eyes firmly, hold down the printed circle of container with your thumbs and push slowly to apply the medicine. Make sure that the dropper tip does not touch eye or eyelashes. After applying, close your eyes for 1 to 5 minutes, pressing the inner corner of your eyes. (If you need to use other eye drops at the same time, wait more than 5 minutes.)
- You can not apply this medicine unless you open the inner plug.
Please open the inner plug in the following procedure before you use a new one.
=How to open the inner plug=
Close the cap, and hold the bottle with both your hands so that the cap faces upwards.
Push the circle printed on the center of the label strongly only once with your thumbs.
- If you miss a dose, apply one dose at the time you notice it. Apply generally from the next time. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include itchy/stinging/bloodshot eyes and blepharitis (erosion of eyelids). If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store at the room temperature (1 to 30 degrees Celsius) away from direct sunlight.
- Discard the remainder. Do not store them. Do not use the medicine that has been stored more than 4 weeks after opening.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.