

# Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 11/2013

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**Juvela N Capsules 100mg

**Active ingredient:**Tocopherol nicotinate

**Dosage form:**opaque red and white capsule (length 16.2mm)

**Imprint or print on wrapping:**(Face) ユベラ N カプセル 100mg, EN100E  
(Back) ユベラ N, 100, JuvelaN, EN100E



## Effects of this medicine

This medicine is effective in improving blood flow in peripheral vessel and lowering cholesterol to improve symptoms such as hypertension.

It is usually used in the treatment of accompanying symptoms resulting from hypertension, hyperlipidemia, and peripheral circulatory disturbance resulting from arteriosclerosis obliterans.

## The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, for adults, take 1 to 2 capsules (100 to 200 mg of the active ingredient) at a time, 3 times daily. The dosage may be adjusted according to the disease, age or symptoms. Strictly follow the instructions.
- If you miss a dose, take a dose as soon as possible when you remember that you missed a dose. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should not take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

## Precautions while taking this medicine

### Possible adverse reactions to this medicine

The most commonly reported adverse reactions include rash, diarrhea, anorexia, and gastric discomfort. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

No pertinent entries.

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

### Storage conditions and other information

- Keep the medicine out of the reach of children. Store it away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

### For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.