

Drug Information Sheet("Kusuri-no-Shiori")

External

Published: 03/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:LAMISIL Cream 1%

Active ingredient:Terbinafine hydrochloride

Dosage form:white cream

Imprint or print on wrapping:



Effects of this medicine

This medicine suppresses fungal proliferation and shows fungicidal effects by inhibiting the enzymes needed for biosynthesis of cellular membrane components of fungi (mold), such as dermatophytes.

It is usually used to treat dermatomycoses such as tinea (athlete's foot) and candidiasis of skin.

The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply to the affected area once a day. Strictly follow the instructions.
- Do not apply the medicine to your eyes. If the medicine gets into your eyes, wash it off with water and consult your doctor or pharmacist since it may cause irritative symptoms.
- If you miss a dose, apply the missed dose when you remember. You should not apply two doses at one time.
- If you accidentally use more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop using this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include local contact dermatitis (rash), itch, redness, erythema, irritation, scale (like scurf), exfoliation (desquamated skin) and skin fissures. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from heat.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.