

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 02/2022

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: Lacrimin ophthalmic solution 0.05%

Active ingredient: Oxybuprocaine hydrochloride

Dosage form: limpid and colorless, ophthalmic solution, 5mL, (bottle) clear and colorless, (cap) very pale blue green

Imprint or print on wrapping:



Effects of this medicine

This medicine blocks the sensory nerve of the conjunctiva and the cornea or the reflex arc of trigeminal nerve which controls tear secretion temporarily in order to suppress excessive tear secretion caused by stimulation. It is usually used to treat secretory epiphora.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods. If you wear contact lenses.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply 1 to 2 drops at a time, 2 to 5 times a day. Strictly follow the instructions.
- Before applying, wash your hands thoroughly with soap and running water. Pull down your lower eyelid lightly and apply the medicine. Be careful not to touch the dropper tip to your eyelid, eyelashes, or eye. After applying, keep your eyes closed without blinking for a while (1 to 5 minutes). Wipe off any excess liquid gently with clean gauze or a tissue. If you need to apply other eye drops at the same time, wait more than 5 minutes.
- If you wear soft contact lenses, take off the lenses and then apply. After an interval of 5 to 10 minutes to wear contact lenses again.
- If you miss a dose, apply the missed dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include corneal disorder (eye pain, focus disorder, foreign body sensation in eyes), blepharitis (erosion of eyelids/redden eyelids), and itchy eyes. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- facial pallor, cold sweat, dizziness on standing up [shock]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of reach of children. Store away from direct sunlight and heat.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.