

Drug Information Sheet("Kusuri-no-Shiori")

External
Revised: 12/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Lulicon Cream 1%

Active ingredient:Luliconazole

Dosage form:white cream

Imprint or print on wrapping:ルリコンクリーム 1%, LULICON, 10g



Effects of this medicine

This medicine shows antifungal action by inhibiting biosynthesis of ergosterol which is a component of fungal cell membrane.

It is usually used to treat dermatomycosis [tinea (tinea pedis, tinea corporis, tinea cruris), candidiasis (erosion interdigitalis, intertrigo) and chromophytosis].

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply to the affected area once a day. Strictly follow the instructions.
- Not for ophthalmic use (cornea or conjunctiva).
- Do not use on marked erosion.
- If you miss a dose, apply a dose as soon as possible when you noticed. If it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally use more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop using this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- Pay attention, this medicine may cause yellowing of your clothes.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include itch, redness, stimulating feeling, contact dermatitis, pain and eczema. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from light, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.