

Drug Information Sheet("Kusuri-no-Shiori")

Internal
Revised: 05/2018

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: MOVER Tablets 100mg**Active ingredient:** Actarit**Dosage form:** white tablets, diameter: 7.6 mm, thickness: 4.1 mm**Imprint or print on wrapping:** モーバー100mg, MOVER100mg, MKC101**Effects of this medicine**

This medicine exerts antirheumatic action mainly by inhibiting delayed allergic reaction and by suppressing cytokine productions. It alleviates joint pain, swelling and stiffness of the chronic rheumatoid arthritis by controlling the pathology of the disease.

It is usually used to treat rheumatoid arthritis.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant, possibly pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 1 tablet (100 mg of the active ingredient) at a time, 3 times a day. Strictly follow the instructions.
- If you miss a dose, take the dose as soon as possible. If it is close to your next dose, skip the missed dose and continue your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine**Possible adverse reactions to this medicine**

The most commonly reported adverse reactions include rash, itching, abdominal pain and hematuria. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- decreased urine output, general edema [nephrotic syndrome]
- fever, dry cough, respiratory distress [interstitial pneumonia]
- gum bleeding or nose bleeding, sore throat, fever [aplastic anemia, pancytopenia, agranulocytosis, thrombocytopenia]
- general malaise, loss of appetite, yellowing of the skin and the white of eyes [liver dysfunction]
- abdominal and back pain, nausea, bloody diarrhea [peptic ulcer, hemorrhagic colitis]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.