# Drug Information Sheet("Kusuri-no-Shiori")

External

Published: 02/2010

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

#### **Brand name: Mycospor Cream 1%**

Active ingredient:Bifonazole

Dosage form: white cream, (tube) pale purple, (cap) purple

Print on wrapping:マイコスポールクリーム

# Effects of this medicine

This medicine shows antimycotic effects by damaging the structure and function of the cellular membrane of the mold (fungi) parasitizing the skin.

It is usually used to treat dermatomycosis [ringworm, candidiasis and tinea versicolor (athlete's foot, tinea, etc.)].

#### Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- · If you are pregnant or breastfeeding.
- •If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines as well as other prescription medicines.)

#### Dosing schedule (How to take this medicine)

Your dosing schedule prescribed by your doctor is((

to be written by a healthcare professional)

- •In general, clean the affected area and then apply this medicine once a day. Strictly follow the instructions. This medicine is effective when applied after a bath.
- •This medicine is only intended for external use. Do not take it orally or apply it to your eyes.
- If you miss a dose, apply the missed dose when you remember. You should not use two doses at one time.
- · If you accidentally use more than your prescribed dose, consult with your doctor or pharmacist.
- •Do not stop using this medicine unless your doctor instructs you to do so.

#### Precautions while taking this medicine

#### Possible adverse reactions to this medicine

The most commonly reported adverse reactions include irritation, dermatitis, redness/erythema, fissures, dander (peeling off like scales), itch, erosion and dryness. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

### Storage conditions and other information

- · Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- •Discard the remainder. Do not store them.

## For healthcare professional use only

For further information, talk to your doctor or pharmacist.