

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 05/2022

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: NATRIX Tablets2

Active ingredient: Indapamide

Dosage form: pale pink tablet (diameter : approx. 6.6mm, thickness : approx. 3.8mm)

Imprint or print on wrapping: ナトリックス 2, KYO252, NATRIX 2, ナトリックス, 2



Effects of this medicine

This medicine reduces fluid volume by increasing urinary excretion and consequently decreases blood pressure. It is usually used to treat essential hypertension.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash etc.) to any medicines or foods.
If you have: anuria, acute renal failure, low sodium/potassium blood levels.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 1 tablet at a time (2 mg of the active ingredient) once a day after breakfast. The dosage may be adjusted according to the age or symptoms. Strictly follow the instructions.
- If you miss a dose, take the missed dose as soon as possible. However, if the next dose is less than 8 hours, skip the missed dose and continue your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking the medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- The medicine may cause dizziness or lightheadedness due to decreased blood pressure. Pay attention when you work at heights, drive a car or operate dangerous machinery.
- Follow your doctor's instructions and take regular blood tests.
- Pay attention that alcohol may intensify the effect of this medicine.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include dizziness, nausea, weakness/malaise, headache and rash. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- widespread erythema, fever, circular/oval shaped red rash [toxic epidermal necrolysis, oculomucocutaneous syndrome, erythema exsudativum multiforme]
- malaise, loss of appetite, nausea [hyponatremia]
- malaise, weakness, arrhythmia [hypokalemia]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store at room temperature (1 to 30 degrees Celsius) away from direct sunlight and moisture.
- Discard the remainder. Do not store them. If you do not know how to dispose of it, please consult the pharmacy or medical institution where you received it. Please do not give it to others.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.