

# Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 08/2023

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:** NIFLAN OPHTHALMIC SOLUTION 0.1%

**Active ingredient:** Pranoprofen

**Dosage form:** limpid and colorless ophthalmic solution, (volume)5mL

**Imprint or print on wrapping:**



## Effects of this medicine

This medicine suppresses inflammation by inhibiting prostaglandin formation which is the causal substance of inflammation.

It is usually used for reduction of eye inflammation such as blepharitis, conjunctivitis, keratitis, and after eye surgery.

## The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.  
If you have: inflammation from eye infection.  
If you wear contact lenses.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is( ( to be written by a healthcare professional))
- In general, apply 1 to 2 drops at a time, 4 times a day. The dosage may be adjusted according to the symptoms. Strictly follow the instructions.
- Before applying, wash your hands with soap and water. Apply the medicine so the tip of eye dropper does not touch the surface of your eye. After applying, close your eyes for 1 to 5 minutes, pressing gently on the inner corner of the eye with your finger. Wipe off any excess liquid with a clean gauze or tissue immediately.
- If you need to apply other eye drops at the same time, wait more than 5 minutes.
- If you miss a dose, apply a dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally use more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop using this medicine unless your doctor instructs you to do so.

## Precautions while taking this medicine

- If you find out you become pregnant, consult with your doctor.
- If you are going start breastfeeding, consult with your doctor.

## Possible adverse reactions to this medicine

The most commonly reported adverse reactions include stinging eyes, conjunctival hyperemia, itchy eyes and blepharitis (eyelid inflammation). If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

No pertinent entries.

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

## Storage conditions and other information

- Keep out of reach of children. Put in the attached accompanying bag, away from direct sunlight and store at room temperature (1 to 30°C).
- Discard the remainder. Do not store them. If you do not know how to discard of it, please ask your pharmacy or medical institution on how to discard them. Do not give it to others.
- To avoid improper use and quality change, do not transfer this solution to any other bottle or container.
- Do not use any other solution in this bottle.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.