

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Published: 01/2019

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: PYDOXAL Tablet 30mg

Active ingredient: Pyridoxal phosphate hydrate

Dosage form: white tablet (diamater: 9.1 mm, thickness: 4.8 mm)

Imprint or print on wrapping: ピドキサール 30



Effects of this medicine

This active vitamin B₆ preparation works on various metabolism in the body as a coenzyme. It is usually used to treat anemia and skin/mucosa/nerve inflammation caused by the lack of vitamin B₆ as well as to supply vitamin B₆ when its intake from meal is deficient.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, take 10 to 60 mg of the active ingredient in 1 to 3 divided doses a day. The dosage may be adjusted according to the age and symptoms. 1 tablet contains 30 mg of the active ingredient. Strictly follow the instructions of your doctor/pharmacist.
- This medicine is designed not to melt until it reaches the intestines. It is to be swallowed whole, not chewed.
- If you miss a dose, take the missed dose as soon as possible and continue your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include hypersensitivity symptoms such as rash. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- muscle pain, numbness in the limbs, stiffness, lassitude, red-brown urine [rhabdomyolysis]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store at room temperature (1 to 30 degrees Celsius), away from direct sunlight and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.