

# Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 06/2021

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:Stomin A Combination Tablets**

**Active ingredient:**Nicotinamide

Papaverine hydrochloride

**Dosage form:**white tablet, diameter: 7.0 mm, thickness: 2.5 mm

**Imprint or print on wrapping:**ストミン A 配合錠, Stomin A



**Effects of this medicine**

This medicine increases the blood flow of the inner ear and prevents the electrolytic fluctuation of inner ear due to noise stimulus.

It is usually used to relieve the symptoms of tinnitus due to inner ear and central neurological disorder.

**The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.**

- If you have previously experienced any allergic reaction (itch, rash etc.) to any medicines or foods.  
If you have glaucoma or cardiac disease.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

**Dosing schedule (How to take this medicine)**

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, for adults, take 2 tablets at a time, 3 times a day after each meal.  
The dose may be adjusted according to your age and symptoms. Strictly follow the instructions.
- If you miss a dose, take a dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should not take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

**Precautions while taking this medicine**

**Possible adverse reactions to this medicine**

The most commonly reported adverse reactions include rash, palpitation, facial flush, constipation, heartburn, headache and dizziness. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

No pertinent entries.

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

**Storage conditions and other information**

- Keep out the reach of children. Store away from light, heat and moisture.
- Discard the remainder. Do not store them.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.