

# Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 05/2020

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:** TRANSAMIN TABLETS 500mg

**Active ingredient:** Tranexamic acid

**Dosage form:** tablet (white-pale yellowish white), (major axis: 17.8mm, minor axis: 7.2mm, thickness: 5.0mm)

**Print on wrapping:** トランサミン 500mg, Transamin500mg, D608



## Effects of this medicine

This medicine exhibits anti-hemorrhagic/anti-allergic/anti-inflammatory effects by suppressing the activity of plasmin engaged in various bleeding symptoms.

It is usually used to treat symptoms as follows: bleeding tendency considered to be due to systemic hyperfibrinolysis, abnormal bleeding considered to be due to local hyperfibrinolysis, erythema/swelling/itch observed in eczema/hives/drug rash/toxicoderma, sore throat/redness/congestion/swelling observed in tonsillitis/pharyngolaryngitis, pain inside the mouth in stomatitis, aphtha (ulcer) in oral mucous membrane.

## Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, for adults, take 750 to 2,000 mg of the active ingredient in 3 to 4 divided doses a day. The dose may be adjusted according to your age and symptoms. This preparation contains 500 mg of the active ingredient in a tablet. Strictly follow the instructions of your doctor and pharmacist.
- If you miss a dose, take the missed dose as soon as possible. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

## Precautions while taking this medicine

### Possible adverse reactions to this medicine

The most commonly reported adverse reactions include loss of appetite, nausea, vomiting, heartburn, itch and rash. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

- spasmodically contracting muscles [convulsion]

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

### Storage conditions and other information

- Keep out the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

### For healthcare professional use only

For further information, talk to your doctor or pharmacist.