

Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 01/2023

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:ZEFNART CREAM 2%

Active ingredient:Liranaftate

Dosage form:white cream, 10g

Imprint or print on wrapping:ゼフナートクリーム 2%、ZY103



Effects of this medicine

This medicine shows antifungal action by inhibiting biosynthesis of the constituent of fungus cells.

It is usually used to treat tinea (athlete's foot, tinea cruris).

The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply this medicine to the affected area once a day. Strictly follow the instructions.
- Do not apply it to eyes (cornea, conjunctiva). If it accidentally gets into your eyes, wash it out with water (If there is still something wrong, consult with your doctor).
- Do not apply it to the severe erosion.
- If you miss a dose, apply the missed dose when you remember. If it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include contact dermatitis (erosion) and dermatitis symptoms such as pruritus, redness, erythema, irritating sensation. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of the reach of children. Store at a room temperature (1 to 30 degrees Celcius), away from direct sunlight and moisture.
- Discard the remainder. Do not store them.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.