

Contains carfentrazone-ethyl, the active ingredient used in Aim® EC.



HERBICIDE

ANTIK™ EC



ON-CONTACT CONTROL WITH LOW USE RATES

Gain higher efficiency and savings in your crops with Antik™ EC, a fast-acting, low use rate herbicide for control of broadleaf weeds. Containing the active ingredient carfentrazone-ethyl, Antik EC is an excellent tank-mix partner approved for use on a broad range of row and specialty crops with no-till burndown requirements.

Antik EC works on contact and offers growers a more efficient way to harvest through its flexible application methods, from pre-plant, postemergence, to harvest. Take control of tough kochia, waterhemp, lambsquarters, and Russian thistle with ease in cotton, rice, corn, and more with the versatile Antik EC.

KEY BENEFITS

- Controls a wide range of tough broadleaves
- Formulated for low use rates
- Excellent tank-mix partner for no till-burndown
- Flexible application methods

KEY USES

- Corn
- Cotton
- Dry Legumes
- Potatoes
- Rice
- Soybeans
- Strawberries
- Tomatoes

PRODUCT NOTES

EPA REGISTRATION NUMBER
91234-247

ACTIVE INGREDIENT
Carfentrazone-ethyl 22.3%

FORMULATION
Emulsifiable Concentrate

HRAC NUMBER
14

SIGNAL WORD
Caution

PACKAGE SIZE
6 x 1 qt

RESTRICTED USE
No

Refer to label for specific use restrictions.



www.AtticusLLC.com



PRODUCT INFORMATION

Antik EC is to be mixed with water, liquid fertilizer or mixtures of water and liquid fertilizer and adjuvants and applied to labeled crops and non-crop areas for selective postemergence control of broadleaf weeds, for sucker control, for burndown prior to planting, as a harvest aid and to defoliate/desiccate labeled crops. Weed control is optimized when the product is applied to actively growing weeds. Antik EC is a contact herbicide. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation.

Extremes in environmental conditions including temperature, moisture, soil conditions, and cultural practices may affect the activity of Antik EC symptoms may be accelerated under moist conditions. Weed control may be reduced when weeds are hardened off by drought and become less susceptible to Antik EC.

Antik EC is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications must not be made within 6 to 8 hours of either rain or irrigation or when heavy dew is present on the crop. Environmental conditions and certain spray tank additives may increase herbicidal symptoms on the crop.

TANK MIXTURES

Antik EC may be tank-mixed with other registered herbicides for controlling a broader spectrum of weeds. Refer to each product labels for mixing instructions, precautions, and restrictions. Follow the most restrictive instructions for each tank mix partner.

When preparing a new tank mix conduct an appropriate compatibility test by mixing proportional amounts of all spray ingredients in a test vessel (jar) prior to tank mixing with other products. Shake the mixture vigorously and allow it to stand for five to ten minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and must not be applied.

Provided the jar test indicates the mixture to be compatible, prepare the tank mixture as follows: Fill the tank one fourth full with water. With the agitator operating, add the labeled amounts of ingredients using the following order: dry granules first and liquid suspensions (flowables) second. As the agitation continues and the tank is filled with water add emulsifiable concentrate products third followed by the addition of water soluble products.

ADJUVANT USE REQUIREMENTS

The use of a quality spray adjuvant is required for optimum performance. Refer to the individual crop sections of this label for specific adjuvant type and use rates.

Refer to product label for complete application and mixing instructions.

©2022 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Antik™ EC is not manufactured, or distributed by FMC, seller of Aim® EC. Aim® is a registered trademark of FMC Corporation or an affiliate. Antik™ is a trademark of Atticus, LLC.

KEY WEEDS

Amaranth; palmer, spiny
Anoda, spurred
Bedstraw, catchweed
Bindweed, field
Buffalobur
Burclover
Carpetweed
Cheeseweed
Cocklebur
Copperleaf, hophornbeam
Corn Spurry
Cotton; GMO varieties, volunteer
Dayflower
Eclipta
Fiddleneck; coast
Filaree; broadleaf, white, redstem
Flixweed
Groundcherry; smooth, Wright's
Jimsonweed
Kochia
Lambsquarters, common
Lettuce, prickly
Mallow; common, Venice
Meadowfoam
Morningglory; entireleaf, ivyleaf, pitted, scarlet
Nettle, burning
Nightshade; American black, black,
Pennycress, field
Pigweed; prostrate, smooth, tumble
Purslane, common
Redmaids
Rocket, London
Sage, lanceleaf
Sesbania, hemp
Shepherdspurse
Smartweed, PA
Sowthistle
Speedwell, Virginia
Spiderwort, tropical
Spurge, prostrate*
Tansymustard
Thistle, Russian
Velvetleaf*
Wallflower, bushy
Waterhemp; common, tall

**Not for Use in California
(Refer to product label for complete list)*

