HERBICIDE



SETTING YOUR SOYBEAN CROP UP FOR SUCCESS.

Maximum yield is the first goal in soybean production so your first step should be to protect your crop from broadleaf and grass weeds with Aquesta[™] FR. Aquesta FR contains sulfentrazone and cloransulam-methyl, the active ingredients found in Authority[®] First DF and in Sonic[®]. This powerful product provides preemergent protection by infiltrating the roots and shoots of cocklebur, green foxtail, ragweed, wild mustard, and many other weeds.

Aquesta FR can be used alone or tank mixed with fertilizers, insecticides, or other herbicides for maximum flexibility. Aquesta FR is adaptable to all tillage systems and can be used as part of a spring burndown program when tank mixed with a compatible burndown herbicide. Adequate rainfall of 1/2-1" is needed for herbicide activation, and once activated, Aquesta FR helps to control many of the weeds that will rob you of the yield you work hard to achieve.

KEY BENEFITS

- Preemergence protection against broadleaf weeds
- Prevention of weed resistance through dual modes of action
- · Can be used in no-till, minimum-till, and conventional tillage systems

KEY USES

Soybeans



PRODUCT NOTES

EPA REGISTRATION NUMBER 91234-41

ACTIVE INGREDIENTS Sulfentrazone 62.1% Cloransulam-methyl 7.9%

FORMULATION Water Dispersible Granule

HRAC NUMBERS 14 2

SIGNAL WORD Caution

PACKAGE SIZE 2 x 10 lb

RESTRICTED USE No

Refer to label for specific use restrictions.



KEY WEEDS



APPLICATION INFORMATION

- Use a standard low pressure herbicide boom sprayer equipped with suitable nozzles and screens. Apply uniformly using properly calibrated nozzles (10 to 40 psi) and screens and strainers no finer than 50 mesh.
- Use 10 to 40 gallons of spray solution per acre. Do not exceed 40 psi spray pressure unless required by the spray nozzle manufacturer.
- Continuous agitation during application is required. Avoid swath overlaps. Shut off spray booms while turning, slowing or stopping, as over application may result. Do not allow Aquesta FR spray mixtures to sit overnight as settling of product and difficulty of re-suspending may occur.
- Avoid all direct, and/or indirect spray contact with non-target plants. Do not apply near desirable vegetation. Allow adequate distance between target area and desirable plants to minimize exposure.

BEST PRACTICES:

- Preemergence and preplant incorporated applications of Aquesta FR require rainfall or irrigation to activate the herbicide.
- The amount of rainfall or irrigation required for activation following application depends on existing soil moisture, organic matter content, and soil texture.
- Observe all instructions, crop restrictions, mixing directions, application precautions, replanting directions, rotational crop guidelines, and other label information of each product when tank mixing Aquesta FR.

Barnyardgrass Cocklebur **Common Sunflower** Green foxtail Kochia Lambsquarters Marestail Moringglory species Nightshade species Palmer Amaranth Pigweed Ragweed Russian Thistle **Smartweed Species** Velvetleaf Venice Mallow Waterhemp Wild Mustard Yellow foxtail

(Refer to product label for complete list)

Refer to product label for complete application and mixing instructions.

©2019 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Aquesta™ FR is not manufactured, or distributed by FMC Corporation, seller of Authority[®] First DF, or Dow Agrosciences, seller of Sonic[®]. Authority[®] First DF is a registered trademark of FMC Corporation. Sonic[®] is a registered trademark of Dow Agrosciences. Aquesta[™] FR is trademark of Atticus, LLC.

