

# FLUAZINAM 40SC SELECT

### Broad spectrum turf fungicide

**ACTIVE INGREDIENT** Fluazinam – *Group 29* 

**% ACTIVE INGREDIENT** Fluazinam . . . . . . . . . . . . 40%

**FORMULATION**Suspension concentrate

**CHEMICAL FAMILY** 2,6-dinitroanilines

BRAND ALTERNATIVE Secure®

**MODE OF ACTION**Oxidative phosphorylation uncoupler

**RESTRICTED USE**No

SIGNAL WORD
Caution

**PACKAGING** 4 x 0.5 gal, 2 x 2.5 gal

**LABELED USES**Golf course tees, greens, fairways and roughs



Fluazinam 40SC Select™ is a broad spectrum, multi-site fungicide to control diseases in turf grasses on golf courses only. The active ingredient, fluazinam, is the only registered fungicide for turf in FRAC 29 group. It has no known resistance and as a multi-site contact, it has a very low risk of developing resistance. When partnered with chlorothalonil-based fungicides, Fluazinam 40SC Select™ provides season-long protection against the toughest turf diseases without interruptions.

#### **PRODUCT BENEFITS**

- Unique multi-site contact active ingredient, reduces resistance management
- Excellent control of dollar spot and 10 additional turf diseases
- Perfect tank mix partner with systemics for inside out protection
- Controls dollar spot that is resistant to other chemistries



## FLUAZINAM 40SC SELECT

Broad spectrum turf fungicide

#### APPLICATION INFORMATION

Apply Fluazinam 40SC Select™ in sufficient water to obtain adequate uniform coverage of the turf. Apply in a minimum spray volume of 40 gal of water per acre on fairways and 90 gal of water per acre on tees and greens.

- Apply at 0.5 fl. oz. per 1,000 sq. ft. (6 acres per gallon) every 14 days.
- Apply after mowing with sufficient water to uniformly cover the turf and allow area to dry completely before irrigation.
- Refer to the product label for additional information about this product and tank mixing.
- Not for use on home lawns or any residential turf.

#### **Turfgrass disease control**

- Use Fluazinam 40SC Select<sup>™</sup> in a preventative disease control program.
- Apply in sufficient water to ensure thorough coverage.
- Apply after mowing or allow sprayed area to completely dry before mowing.
- For control of foliar diseases, allow sprayed area to completely dry before irrigation.
- For optimum turf quality and disease control, use Fluazinam 40SC Select™ in conjunction with turf management practices that promote good plant health and optimum disease control.
- Evaluate spray additives prior to use. Label directions are based on data obtained with no additives.
- Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.





#### Restrictions

- DO NOT exceed 258 fl oz of Fluazinam 40SC Select™ (8.4 pounds active ingredient) per acre, per year.
- DO NOT apply more than twelve (12) applications at 21.5 fl oz per acre (0.5 fl oz/1000 sq ft), per year.

#### Labeled diseases

Algal scum Gray snow mold Pink patch

Anthracnose Large patch Pink snow mold

Brown blight Leaf spot Red thread
Brown patch Melting out Stem rust

Dollar spot

Read label for full application and rate information.