

Phostrol®

AGRICULTURAL FUNGICIDE

SYSTEMIC ACTIVITY FOR EFFECTIVE DISEASE CONTROL.

Phostrol® provides both preventive and curative disease control across 16 horticultural crops. Phostrol is a phosphite fungicide that delivers proven, cost-effective disease control for improved crop quality and higher yield potential against a host of diseases. Phostrol is an excellent fit for your integrated disease management program.

COMPETITIVE ADVANTAGES

PREVENTION AND CONTROL

- Fire and late blights, canker, downy mildew, Phytophthora spp., storage rot diseases, pecan scab, silver scurf

PRODUCT ATTRIBUTES

- No odor: clear, virtually odorless formulation
- Neutral pH: requires no buffering for tank-mix compatibility and the integration with most fungicides
- Foliar application flexibility: by air, ground or chemigation

KEY CROP USES

- Brassica (cole), caneberry, cucurbits, citrus, grapes, leafy vegetables, legume vegetables, pome and stone fruit, onions, potato, strawberry, tobacco, tomato, tree nuts and pistachios, tuberous and corm vegetables

AT-A-GLANCE INFO

EPA REG. NO. 55146-83

ACTIVE INGR. mono- and dibasic sodium, potassium and ammonium phosphites (53.6%)

FORMULATION liquid

CHEM. FAMILY phosphite

GROUP NO. 33

MODE OF ACT. unknown

REI 4 hours

SIGNAL WORD caution

RESTRICTED no

PACKAGE SIZE 2.5 gal container (2 per case), 265 gal tote

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

Agri-Mycin® 50

ChampION™

Mycoshield®

 **Nufarm**
Grow a better tomorrow

Phostrol®

SYSTEMIC ACTIVITY FOR EFFECTIVE DISEASE CONTROL.

KEY USES

Asparagus
Avocado
Blueberries
Citrus
Caneberries
Cranberries
Cucurbit crops
Ginseng
Grapes
Hops
Leafy vegetables
Legume vegetables
Tree nuts and Pistachios
Onion
Potatoes
Strawberries
Tomatoes

KEY DISEASES CONTROLLED

(See product label for complete list)

FOLIAR APPLICATION:

Canker
Downy mildew
Fire blight
Late blight
Phytophthora spp. root rot
Pecan scab
Red stele
Storage rot diseases

IN-FURROW APPLICATION:

Storage rot diseases
Pink rot
Pythium leak

PAINT OR SPRAY APPLICATION:

Canker

POST-HARVEST (POTATOES ONLY):

Late blight
Pink rot
Silver scurf

USE SUMMARY

RATE	CROP
FOLIAR APPLICATION RECOMMENDATIONS	
2.0 - 4.0 pints/A	Peppers
2.0 - 5.0 pints/A	Legume Vegetables; All members of Crop Group 6
2.5 pints/A	Hops
2.5 - 3.0 pints/A	Pineapple (see label for dip application)
2.5 - 3.75 pints/A	Onion (dry bulb)
2.5 - 5.0 pints/A	Asparagus
	Avocado
	Blueberry
	Brassica (Cole); All members of Crop Group 5
	Citrus; All members of Crop Group 10
	Cucurbits; All members of Crop Group 9
	Grapes
	Leafy Vegetables; All members of Crop Group 4
	Pome Fruit; All members of Crop Group 11
	Stone Fruit; All members of Crop Group 12
	Strawberry (see label for dip application)
	Tobacco ¹
	Tomato
	Tree Nuts ² and Pistachio; All members of Crop Group 14 and Pistachios
2.5 - 10.0 pints/A	Tuberous and Corm Vegetables; All members of Crop Group 1C and 1D
4.5 pints/A	Caneberry ¹ ; Blackberry, Loganberry, Red and Black Raspberry, Hybrids, etc.
	Ginseng ¹
5.0 - 6.0 pints/A	Cranberry
IN-FURROW APPLICATION RECOMMENDATIONS	
3.75 - 10.0 pints/A	Tuberous and Corm Vegetables; All members of Crop Group 1C and 1D

¹ Not registered for use in California.

² Not registered for use in California for treatment of pecan scab.

POST HARVEST – POTATOES ONLY

Apply 0.1 gallon of Phostrol® per ton of tubers in 0.5 gallon of water as spray. Ensure complete and even coverage.

See label for complete directions for use.



nufarm.com/uscrop

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/uscrop to download a full label.

©2020 Nufarm. Important: Always read and follow label instructions. Agri-Mycin®, ChampION™, Mycoshield® and Phostrol® are trademarks of Nufarm.



15-AG-0033-C