

EASY-TO-MIX LIQUID FORMULA AVAILABLE THROUGH GOWAN USA

Key Features and Benefits

- **Fast Knockdown** of difficult to control mites and whiteflies
- **Lasting control** - 30 to 45 days residual
- **Long list of tolerant plant species** - even on bracted poinsettia
- **Effective against most major mite species** including:
 - Twospotted Spider Mite
 - Broad Mite
 - European and Southern Red Mite
 - Rust and Bud Mite
- **Effective against greenhouse and sweet potato whitefly**, including Q-biotype in *Bemisia tabaci* populations.
- **Controls leafhopper and certain aphids** as listed on the label

PRODUCT PROFILE

A.I. Load	42.47% Pyridaben
Formulation:	SC
Signal Word:	CAUTION
IRAC #:	21A
Use Rate:	6.5 - 9.5 oz per acre OR 0.15 - 0.22 oz per 1000 sq ft



GENERAL INFORMATION

800.883.1844

www.gowanco.com






SANMITE® SC IS A HIGHLY EFFECTIVE MITICIDE/INSECTICIDE THAT CONTROLS MITES AND WHITEFLIES ON THE FOLLOWING CROPS:

- Ornamental plants
- Flowers and Foliage Plants
 - Greenhouse tomatoes
 - Greenhouse cucumbers


Directions For Use:

1. Apply in sufficient water to ensure thorough coverage to foliage. Thorough coverage is required for optimal control.
2. In general, no additives are needed, but certain oils and spray adjuvants have been shown to improve performance. Contact your distributor sales representative or Gowan USA representative for details.
3. Use a minimum of 6.5 ounces per acre.
4. Treat plants when pests are immature or at a susceptible stage. For best results, Sanmite SC should be applied as populations build, prior to reaching economic threshold.
5. Use 100-400 gallons of spray solution per broadcast acre for optimal performance.

**Whitefly Nymphs per 10 cm²
 (5 cm² from 2 plants per plot)**



**Whitefly Nymphs per 10 cm²
 (5 cm² from 2 plants per plot)**



RATE COMPARISON

SANMITE[®] SC MITICIDE/INSECTICIDE	7 floz.	8 floz.	9.5 floz.
SANMITE MITICIDE	4.4 oz.	5 oz.	5.94 oz.