

GROUP	4	14	HERBICIDE
-------	---	----	-----------



Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	9.02%
Dichlorprop-p, 2-ethylhexyl ester	5.19%
Dicamba, acid	0.59%
Carfentrazone-ethyl	0.47%
OTHER INGREDIENTS:	84.73%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.51 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 5.98%.
- 0.30 lb. (R)-2-(2,4-Dichlorophenoxy)propionic acid equivalent per gallon or 3.51%.
- 0.050 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 0.59%.
- 0.040 lb. Ethyl α,2-dichloro-5-[4(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate per gallon or 0.47%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

KEEP FROM FREEZING



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

- All mixers, loaders, applicators and other handlers must wear:
- long-sleeved shirt and long pants,
 - shoes and socks,
 - waterproof gloves.
 - chemical resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater Advisory: This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls,
- chemical-resistant gloves made of any water-proof material,
- shoes plus socks, and
- protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

1. Product Description

SpeedZone® Southern EW Broadleaf Herbicide for Turf is a selective herbicide for post-emergence control of listed broadleaf weeds commonly found in turfgrasses. It can be used on established warm- and cool-season turfgrasses as listed.

This product offers these advantages:

- Excellent post emergent activity with proven performance for broadleaf weed control in turfgrass.
- Cool weather performance
- High selectivity (turfgrass safety) in established warm- and cool-season turfgrass.
- Provides rapid and effective weed control for common and troublesome weed species in turfgrass, e.g., spurge, pennywort (dollarweed), dandelion, and white clover.
- Weed control is fast with symptoms showing within hours. Generally, the initial visual symptoms can be noticed within hours of the application and plant death can occur within 7 to 14 days.

2. Use Restrictions

- For ground application only; aerial applications are not permitted. Do not apply this product through any type of irrigation system.
- Do not allow livestock to graze on any areas treated with this product.
- Do not apply this product to bare ground.
- Do not formulate this product into other end-use products.

3. Resistance Management

The active ingredients in this product belong to two different herbicide groups: Group 4 (synthetic auxins) and Group 14 (PPO/protox inhibitors). To maintain the long-term effectiveness of herbicides, consult your local extension specialist, certified advisor, and/or manufacturer representative for resistance management strategies recommended in your area.

4. Spray Preparation And Tank Mixes

SpeedZone Southern EW Broadleaf Herbicide for Turf is a stable emulsion in water (EW) that can be diluted with water or liquid fertilizer to form a stable emulsion. Aqueous micro-emulsions are non-flammable and offer good miscibility with water.

Mixing with water:

Add one-half the required amount of water to the spray tank. Then add this product slowly with agitation and complete filling the tank with water. Mix thoroughly and continue agitation while spraying. When this product is left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Do not use tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer the spray solution to alter the pH range as appropriate.

Mixing with liquid fertilizers:

Use suitable sources and rates of fertilizer based upon recommendations of your fertilizer supplier or State Extension Service Specialist.

Always verify physical compatibility with a jar test before large scale mixing. The jar test can be conducted by mixing all components in a small container in proportionate quantities. If the mixture separates after standing and can be mixed readily by shaking, then the mixture can be used and applied with spray equipment providing continuous agitation. If large flakes, sludge, gels, or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer must not be prepared as a tank mixture.

Liquid fertilizers are either solutions (true fluids) or suspensions. Physical compatibility of this product is adequate with liquid nitrogen solutions. Mixing this product with suspensions or N-P-K solutions may not be satisfactory (may be marginal) without pre-mixing this product with water. Pre-mixing this product with 2 parts water will ensure that the emulsifiers are activated enabling the herbicide to be suspended in the fertilizer.

Adjuvants and spray additives:

Adjuvants (such as surfactants, spreaders, spreader-stickers, spray thickeners, foaming agents, activators, detergents, and drift reducing agents) combined with this product can damage the leaf tissue of turfgrass. If any discoloration or cosmetic effects are objectionable or would be unacceptable, then the use of adjuvant(s) would not be recommended. Do not use adjuvants and spray additive tank-mix combinations unless your experience indicates that the tank mixture will not result in objectionable turf injury.

5. Ground Equipment

Spray distribution: The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator. Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate nozzles, tips, and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Use spray volumes of 3 to 175 gallons per acre (9 fl.oz. to 4 gal per 1000 sq.ft.). Use higher spray volumes for dense weed populations.

- Calibration and proper application are essential when using this product.
- Over-application or rates above those specified on this label can cause turf injury.
- Hand-held technique: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. These motions result in uneven coverage. Instead, the nozzle should be held stationary at the proper height.

Hand operated sprayers including backpack sprayers and compression sprayers are appropriate for small turfgrass areas.

Low Volume Spray Application Equipment: Apply 3 to 22 gallons of total spray solution per acre (9 fl.oz. to 2 qt/1000 sq.ft.). Uniformly wet leaf surfaces. Higher spray volumes may be required for dense weed infestations, difficult to control weeds, mature weeds, or during adverse/extreme environmental conditions.

After using this product, clean sprayer with soap or detergent and water, or an approved spray tank cleaner and rinse thoroughly before applying other pesticides.

6. Spray Drift Management**6.1 For Ornamental Turfgrass Sites**

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash from treated areas onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

6.2 For Agricultural and Non-Cropland Sites

When this product is used in "commercial sod production," the following Best Management Practices for reducing spray drift apply.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make

applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Additional requirements for ground boom application: Do not apply with a nozzle height greater than 4 feet above the crop canopy.

7. Where To Use

- **Ornamental Turfgrass sites:**
- **Residential/domestic sites** including areas associated with household or home life including apartment complexes, and condominiums.
- **Ornamental turf sites** including turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, playgrounds, fairgrounds, and athletic fields.
- **Institutional sites** including properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses, and office buildings.
- **Non-cropland sites:** including farmyards, fencerows or fence lines, highway rights-of-way (principal, interstate, county, private, and unpaved roads): Roadsides, roadside ditches, road shoulders, road embankments, dividers, and medians; Industrial sites: Lumberyards, tank farms, fuel or equipment storage areas; Municipal, state, and federal lands: Airports and military installations; railroad rights-of-ways, railroad yards, railroad crossings and railroad bridge abutments; Utility rights-of-way: telephone, pipeline, electrical powerlines, and communication transmission lines.
- **Agricultural site:** Commercial sod production

Prohibitions of Sites:

- Do not apply this product to 'Floritam' and 'Bitterblue' varieties of St. Augustinegrass, bentgrass mowed under 1/4 inch, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, estuaries (salt water bays), or wetlands (swamps, bogs, potholes, or marshes). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to agricultural irrigation water or irrigation ditch banks or canals.
- Do not apply to golf course greens.

8. How Much To Use

Use Rates and Spray Volumes:

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

Table 1. Use Rates For Ornamental Turfgrass, Sod Farms, and Non-Cropland		
Species	Rate	Spray Volume
Kentucky bluegrass, annual bluegrass, annual ryegrass, perennial ryegrass, tall fescue, red or fine leaf fescues, bentgrass mowed at 1/2 inch or higher	4 to 6 pt/acre (1.5 to 2.2 fl.oz./ 1000 sq.ft.)	3 to 175 gal/acre (9 fl.oz. to 4 gal/ 1000 sq.ft.)
Bermudagrass, bahiagrass, buffalograss, centipedegrass, seashore paspalum, St. Augustinegrass (excluding 'Floritam' and 'Bitterblue' varieties), zoysiagrass	2 to 4 pt/acre (0.7 to 1.5 fl.oz./ 1000 sq.ft.)	3 to 175 gal/acre (9 fl.oz. to 4 gal/ 1000 sq.ft.)
<p>Note: It is impossible to test all environmental conditions for the listed turfgrasses. We suggest testing this product on a small area and observe the treated area for 30 days to determine the acceptability of turf discoloration.</p> <p>SpeedZone Southern EW Broadleaf Herbicide for Turf should only be applied to turfgrass species that are listed in Table 1 unless trial use indicates that the turf species not listed is tolerant to this product.</p>		

Limitations on broadcast treatments for ornamental turfgrass, sod farms, and non-cropland:

The maximum application rate is 6 pints of product per acre per application [0.38 lb 2,4-D ae, 0.23 lb dichlorprop ae, 0.038 lb dicamba ae, 0.030 lb carfentrazone per acre per application]. The maximum number of broadcast applications is limited to 2 per year. The minimum interval between applications is 21 days for sod farms and 30 days for non-cropland. The maximum seasonal rate is 12 pints of product per acre [0.77 lb 2,4-D ae, 0.45 lb dichlorprop ae, 0.075 lb dicamba ae, 0.060 lb carfentrazone per acre].

Spot Treatment with Hand Operated Sprayers (including backpack sprayers and pump-up type sprayers):

- Spot treatment cannot exceed more than 1,000 sq.ft. in any given acre.
- Apply any time the emerged broadleaf weeds are actively growing.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Over application or rates above those specified on this label including excessive overlaps of this product can cause turf injury and will result in illegal residues.
- Follow-up applications as spot treatments at a 30 day interval are advised for more mature weeds, for dense infestations, and for adverse environmental conditions.
- **For turfgrasses listed in Table 1 with the 4 to 6 pt/acre rate:** Mix 1.5 to 2.2 fl.oz. of this product per 1 gallon of water for treatment of approximately 1000 sq.ft. of turfgrass.
- **For turfgrasses listed in Table 1 with the 2 to 4 pt/acre rate:** Mix 0.7 to 1.5 fl.oz. of this product per 1 gallon of water for treatment of approximately 1000 sq.ft. of turfgrass.

9. Application Timing

Spring and fall treatments are preferred to summer treatments for older, drought-stressed weeds. Fall applications will control emerged winter annuals and perennials such as henbit, chickweed, clover, and ground ivy.

A second broadcast application or a follow-up spot treatment is suggested for more mature weeds, for dense infestations, and for adverse environmental conditions. Do not make more than 2 broadcast treatments of this product per site per year. Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment.

Timing Factors Which Affect Weed Control

- Weed control is more effective when the daytime air temperature is above 50°F, soil moisture is adequate, and target weeds are young and actively growing.
- Rainfast in as little as 3 hours.
- If dry conditions exist, irrigation 24 hours before and 24 hours after the application will increase weed control.
- Higher spray volumes may increase weed control during adverse conditions.

Timing Factors Which Affect Turfgrass Tolerance

- Turf species listed on this label may exhibit temporary discoloration under adverse environmental conditions.
- Temperatures over 90°F, moist soil, and high humidity will tend to increase herbicide activity. These conditions will also increase the possibility of temporary turf discoloration.

- Other conditions which may increase the possibility of turf injury include: disease, insect, and nematode stress; low light (shaded) areas, low soil pH, improper mowing, or improper applications of fertilizer and pesticides.
- If injury occurs, turf will resume normal color and growth after mowing.

For Newly Seeded Areas:

Delay application of this product to grass seedlings until after the second mowing.

For Newly Sodded, Sprigged, or Plugged Areas:

The application of this product should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Reseeding interval:

Treated areas may be reseeded 1 week after application.

Mowing:

Delay mowing 2 days before and until 2 days after the application of this product.

10. Broadleaf Weeds Controlled

SpeedZone Southern EW Broadleaf Herbicide for Turf will control or suppress the following broadleaf weeds. Apply any time the emerged broadleaf weeds are susceptible.

Broadleaf Weeds		
Aster, white heath & white prairie	Filaree, whitestem & redstem	Poison oak
Bedstraw	Florida betony	Prickly lettuce (*compass plant)
Beggarweed, creeping	Florida pusley	Puncturevine
Bindweed	Ground ivy	Purple cudweed
Black medic	Groundsel	Purslane
Broadleaf plantain	Hawkweed	Ragweed
Buckhorn plantain	Healall	Red sorrel (*sheep sorrel)
Bull thistle	Henbit	Shepherd's purse
Burdock, common	Innocence (Blue-eyed Mary)	Spurge
Buttercup, creeping	Knotweed	Thistle
Carpetweed	Lambsquarters	Virginia buttonweed
Chickweed	Lawn burweed	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Chicory	Lespedeza	Wild carrot
Cinquefoil	Mallow, common	Wild garlic
Clover	Matchweed	Wild geranium
Cudweed	Mouseear chickweed	Wild lettuce
Curly dock	Nettle	Wild mustard
Dandelion	Old world diamond flower	Wild onion
Dayflower	Oxalis (*yellow woodsorrel & creeping woodsorrel)	Wild strawberry
Deadnettle	Parsley-piert	Wild violet***
Dock	Pennsylvania smartweed	Yarrow
Dogfennel	Pepperweed	Yellow rocket
English Daisy	Pigweed	
False dandelion (*spotted catsear & common catsear)	Pineappleweed	
Field bindweed (*mornningglory & creeping jenny)	Plantain	
Field oxeye-daisy (*creeping oxeye)	Poison ivy	

* Synonyms
 *** For best results, apply in the spring when wild violets are blooming or apply a late fall application followed by a spring application.

STORAGE AND DISPOSAL (cont.)

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For Plastic Containers – Nonrefillable with capacities greater than 5 gallons:

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT: Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner of use or application, etc. Such factors and conditions are beyond the control of the manufacturer, and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons:

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(cont. on next column)

PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

SpeedZone® and ®Checkered Flag/Label Design are registered trademarks of PBI-Gordon Corporation.

874/7-2019 AP073019
EPA REG. NO. 2217-1031



MANUFACTURED BY
PBI/GORDON CORPORATION
P.O. BOX 860350
SHAWNEE, KANSAS 66286
PBIGordonTurf.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.