

# TIGRIS™ IMAZETHAPYR 2SL

**HERBICIDE FOR USE IN ALFALFA, CLOVER, PEAS AND BEANS, PEANUTS, AND SOYBEANS**  
**Not for use on CLEARFIELD® rice or any other rice varieties or hybrids.)**

<b>ACTIVE INGREDIENT:</b>	<b>By Wt.</b>
Ammonium salt of imazethapyr ( )-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid*	22.87%
<b>OTHER INGREDIENTS:</b>	77.13%
<b>TOTAL:</b>	100.0%

\*Equivalent to 21.6% ( )-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxylic acid (1 gal. contains 2.0 lbs. of active ingredient as the free acid).

## KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
 (If you do not understand the label, find someone to explain it to you in detail.)

<b>FIRST AID</b>	
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of an emergency involving this product, call CHEMTREC at <b>1-800-424-9300</b> .	

EPA Reg. No.: 93809-3-92647



**Manufactured for: Tigris, LLC**  
 P.O. Box 250  
 10025 Hwy. 264 Alternate  
 Middlesex, NC 27557

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as butyl rubber  $\geq 14$  mils, or natural rubber  $\geq 14$  mils, or neoprene rubber  $\geq 14$  mils, or nitrile rubber  $\geq 14$  mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

**Groundwater Advisory and Proper Handling Instructions**

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Tigris Imazethapyr 2SL** may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes or reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

**DO NOT** apply **Tigris Imazethapyr 2SL** through any type of irrigation system.

**Tigris Imazethapyr 2SL** must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixture.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not mix or allow coming in contact with oxidizing agents. Hazardous chemical reactions may occur.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. This labeling must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations on this label and on the labels of products used in combination with **Tigris Imazethapyr 2SL**. Do not use **Tigris Imazethapyr 2SL** other than in accordance with the instructions set forth on this label. The use of **Tigris Imazethapyr 2SL** not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination.



**Manufactured for: Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as butyl rubber  $\geq 14$  mils, or natural rubber 14 mils, or neoprene rubber 14 mils, or nitrile rubber 14 mils
- Shoes plus socks

### PRODUCT INFORMATION

Not for use on CLEARFIELD® rice or any other rice varieties or hybrids.

**Tigris Imazethapyr 2SL** kills weeds by root and/or foliage uptake and rapid translocation to the growing points. Adequate soil moisture is important for optimum **Tigris Imazethapyr 2SL** activity. When adequate soil moisture is present, **Tigris Imazethapyr 2SL** will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

Occasionally, internode shortening and/or temporary yellowing of crop plants occurs following **Tigris Imazethapyr 2SL** applications. These effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

When organophosphate (such as chlorpyrifos) or carbamate insecticides are tank-mixed with **Tigris Imazethapyr 2SL**, temporary injury may result to the treated crops.

Use of **Tigris Imazethapyr 2SL** herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible. Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall) **Tigris Imazethapyr 2SL** may cause injury to subsequent planted crops. Vegetable crops and particularly sugar beets are sensitive to **Tigris Imazethapyr 2SL** residues in the soil.

Naturally occurring biotypes\* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include the sulfonylureas, the sulfonamides and the pyrimidyl benzoates. If naturally occurring ALS/AHAS resistant biotypes are present in a field, **Tigris Imazethapyr 2SL** and/or any other ALS/AHAS enzyme inhibiting mode of action herbicide must be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

\*A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Replanting: If replanting is necessary in a field previously treated with **Tigris Imazethapyr 2SL**, the field may be replanted to soybeans, or peanuts (imidazolinone resistant corn), lima beans or Southern peas. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of **Tigris Imazethapyr 2SL**.

### EDIBLE LEGUME VEGETABLES

Reduced crop growth, quality, yield and/or delayed maturity may result from **Tigris Imazethapyr 2SL** application to edible legume vegetables. Since crop maturity may be delayed, timing of harvest may need to be adjusted accordingly.

Use **Tigris Imazethapyr 2SL** ONLY if proper agronomic practices have been utilized, including good soil fertility, proper crop rotation, disease and insect management and tillage practices that eliminate compaction and hardpans. Plant peas, lentils or lima beans at least 1/2 inch deep to reduce risk of crop injury.

### USE AREA RESTRICTIONS

In New York State – Not for Sale or Use on Long Island.

### RESTRICTIONS

- DO NOT apply **Tigris Imazethapyr 2SL** if planting is delayed and chance of frost prior to maturity is likely.
- DO NOT apply **Tigris Imazethapyr 2SL** if cold and/or wet conditions are present or predicted to occur within one week of application.
- DO NOT apply **Tigris Imazethapyr 2SL** post-emergence after crop has begun to flower or crop injury may result. (Refer to specific legume vegetable crop for specific application timings listed).



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## MIXING INSTRUCTIONS

**POST-EMERGENCE APPLICATIONS OF TIGRIS IMAZETHAPYR 2SL REQUIRE THE ADDITION OF AN ADJUVANT AND A FERTILIZER SOLUTION.**

**NOTE: DO NOT** use fertilizer solutions in the state of California.

### ADJUVANTS

**CROP OIL CONCENTRATE:** A petroleum or vegetable seed based oil concentrate may be used. Use methylated seed oils are when weeds are under moisture or temperature stress. Use methylated seed oils at the rate of 1.0% v/v (1 gal. per 100 gals. of spray solution), or use a crop oil concentrate at 1.25% v/v (1.25 gals. per 100 gals. of spray solution). **DO NOT** include a CROP OIL CONCENTRATE when applying **Tigris Imazethapyr 2SL** to edible legume vegetable crops.

### OR

**SURFACTANTS:** Use a non-ionic surfactant containing at least 80% active ingredient. Apply the surfactant at the rate of 0.25% v/v (1 quart per 100 gals. of spray solution). An organosilicone surfactant or dry surfactant may be used in place of a non-ionic surfactant.

### AND

**(All States Except California)**

### FERTILIZER SOLUTION

Acceptable nitrogen-based fertilizers include liquid fertilizers (such as 28%N, 32%N, or 10-34-0) may be applied at the rate of 1.25 to 2.5 gals. per 100 gals. of spray solution. Use the higher rate within the specified rate range when weeds are under moisture or temperature stress. Instead of a liquid fertilizer, use spray grade ammonium sulfate at the rate of 12 – 15 lbs. per 100 gals. of spray solution.

**NOTE:** Fertilizer solution is not required in **Tigris Imazethapyr 2SL** applications in use areas south of interstate highway 40, except in the states of Texas, New Mexico and Oklahoma.

Fill the spray tank one-half full with clean water. Use a calibrated measuring device to measure the required amount of **Tigris Imazethapyr 2SL**. Add **Tigris Imazethapyr 2SL** to the spray tank while agitating. Add adjuvants and fill the remainder of the tank with water.

### TANK-MIX COMBINATIONS WITH OTHER HERBICIDES

If other herbicides are tank-mixed with **Tigris Imazethapyr 2SL**, while agitating, add components in the following order:

1. Fill spray tank 1/2 full with clean water.
2. Add soluble packet products and thoroughly mix.
3. Add WP (wetable powder), DG (dispersible granule), DF (dry flowable) or liquid flowable formulations not in soluble packets.
4. Add **Tigris Imazethapyr 2SL** and thoroughly mix.
5. Add other aqueous solution products.
6. Add EC (emulsifiable concentrate) products.
7. Add surfactant or crop oil to the spray tank.
8. Add liquid fertilizer.
9. While agitating, fill the remainder of the tank with water.

To avoid injury to sensitive crops, spray equipment used for **Tigris Imazethapyr 2SL** applications must be drained and thoroughly cleaned with water before being used to apply other products.

When **Tigris Imazethapyr 2SL** is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. **DO NOT** exceed label dosages. **Tigris Imazethapyr 2SL** cannot be mixed with any product containing a label prohibiting such mixtures.

### SPRAYING INSTRUCTIONS

**DO NOT** apply when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include, but are not limited to, leafy vegetables and sugar beets.

### GROUND APPLICATIONS

Uniformly apply with properly calibrated ground equipment in 10 or more gals. of water per acre. Apply with a spray pressure of 20 to 40 psi.

To ensure thorough coverage, use a minimum of 20 gals. of water per acre when applying **Tigris Imazethapyr 2SL** to minimum or no-till crops. Use higher gallonage for fields with dense vegetation or heavy crop residues. Adjust the boom height to ensure proper coverage of weed foliage (according to the manufacturer's recommendation). Use only flat-fan nozzle tips for post-emergence applications.

Avoid overlaps when spraying.

### TIGRIS IMAZETHAPYR 2SL APPLICATIONS WITH A LOW-VOLUME SPRAYER

**Tigris Imazethapyr 2SL** may be applied to soybeans with a low-volume (Spra-Coupe® type) sprayer. When applying **Tigris Imazethapyr 2SL** with a low-volume sprayer, spray the weeds before they reach the maximum size listed in this label. Adequate control of weeds is dependent upon good spray coverage of the weeds. The sprayer must be calibrated to deliver the specified spray volume and pressure to ensure adequate spray coverage of the weeds.

When applying **Tigris Imazethapyr 2SL** with a low-volume sprayer, apply a minimum of 10 gals. per acre of spray solution with a nozzle pressure between 40 – 60 psi for optimum coverage.



**Manufactured for: Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## AERIAL APPLICATIONS

**Tigris Imazethapyr 2SL** herbicide may be applied by air to crops listed in this label unless otherwise noted.

Uniformly apply with properly calibrated aerial equipment in 5 or more gals. of water per acre. When applied POST-EMERGENCE, the addition of a non-ionic surfactant AND fertilizer solution are required for optimum weed control. Apply a non-ionic surfactant at the rate of 1 quart per 100 gals. of spray solution OR a crop oil concentrate at the rate of 1.25 gals. per 100 gals. of spray solution AND a liquid fertilizer at the rate of 1.25 gals. per 100 gals. of spray solution. (See instructions under **APPLICATION INFORMATION – POSTEMERGENCE**).

### Mandatory Spray Drift

#### Aerial Applications

- Do not release spray at a height greater than 10 ft. above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a medium to ultra coarse spray droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

#### Ground Applications

- Apply with a nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a medium to ultra coarse spray droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

#### Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.
- IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

#### Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

#### Controlling Droplet Size - Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

#### BOOM HEIGHT - Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

#### RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aurally to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

#### SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

#### TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

#### TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

#### WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## RESISTANCE MANAGEMENT

For resistance management, **Tigris Imazethapyr 2SL** is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to **Tigris Imazethapyr 2SL** and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same use site. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Avoid the consecutive use of **Tigris Imazethapyr 2SL** or other target site of action Group 4 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Base herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Scout fields prior to application to identify the weed species present and their growth state to determine if the intended application will be effective.
- Scout fields after application to verify that the treatment was effective and to monitor weed populations for early signs of resistance development.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

Suspected herbicide-resistant weeds may be identified by these indicators:

- o Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- o A spreading patch of non-controlled plants of a particular weed species; and
- o Surviving plants mixed with controlled individuals of the same species.

Report any incidence of non-performance of this product against a particular weed species to your Tigris, LLC retailer or representative. If resistance is suspected, treat weed escapes with an herbicide having a different mechanism of action and/or use non-chemicals means to remove escapes, as practical, with the goal of preventing further seed production.

## APPLICATION INFORMATION

### POST-EMERGENCE

**Tigris Imazethapyr 2SL** is effective in controlling weeds in conservation tillage as well as in conventional production systems. Apply **Tigris Imazethapyr 2SL** herbicide as an early post-emergence treatment when weeds are actively growing and before they exceed a height of 3 inches, unless otherwise indicated. Delay application until the majority of the weeds are at the specified growth stage. Base application timing on weed size and not crop growth stage. Apply **Tigris Imazethapyr 2SL** to crops and weeds that are actively growing.

An adjuvant (either a crop oil concentrate or a surfactant) and a nitrogen-based fertilizer must be added to the spray solution for optimum weed control activity. See the **ADJUVANTS** section under **MIXING INSTRUCTIONS** for specific instructions.

When **Tigris Imazethapyr 2SL** is applied post-emergence, absorption will occur through both the roots and foliage. Susceptible weeds stop growing and either die or are not competitive with the crop. **Tigris Imazethapyr 2SL** not only controls many existing broadleaf and grass weeds when applied post-emergence, it also provides control of susceptible weeds that may emerge after application.

For maximum weed control, cultivate 7 – 10 days following a post-emergence **Tigris Imazethapyr 2SL** application. This timely cultivation will enhance residual weed control, especially under dry conditions.

Apply **Tigris Imazethapyr 2SL** a minimum of one hour before rainfall or overhead irrigation.

Unusually cool temperatures (50°F or less) reduce photosynthesis and transpiration and thus reduce uptake, translocation, and efficacy of **Tigris Imazethapyr 2SL** herbicide in weeds. Delaying a **Tigris Imazethapyr 2SL** application for 48 hours from the time the temperature increases above 50°F, if air temperature has been below 50°F for 10 hours or more hours, will improve weed control and reduce crop response.

### NO-TILL/MINIMUM TILLAGE AND DOUBLE CROP SOYBEANS

**Tigris Imazethapyr 2SL** controls existing weeds and provides residual control of most weeds when applied early post-emergence to soybeans in no-till or minimum tillage and double crop soybean production systems. The application may be applied either before or after emergence of the crop. (Refer to the **WEEDS CONTROLLED POST-EMERGENCE** chart for weeds controlled and specified weed size).

If **Tigris Imazethapyr 2SL** is applied prior to emergence of the crop, and weeds exceed the specified size, add a contact herbicide to **Tigris Imazethapyr 2SL** to enhance control. (See instructions for **NO-TILL OR REDUCED TILLAGE** under the **PRE-EMERGENCE** section of this label).

### SOIL APPLICATIONS

**Tigris Imazethapyr 2SL** provides effective weed control in conservation tillage systems designed to meet conservation compliance requirements. **Tigris Imazethapyr 2SL** can be applied as an early pre-plant, pre-plant incorporated, or pre-emergence treatment in soybeans. It can also be applied in conventional, minimum tillage and no-till production systems. The application method of choice will depend on the anticipated weed spectrum and the preference of the applicator.

Adequate soil moisture is required for optimum activity. Rainfall or overhead irrigation is necessary to move **Tigris Imazethapyr 2SL** into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after treatment, cultivate to control escaped weeds. When adequate moisture is received after dry conditions, **Tigris Imazethapyr 2SL** will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

**Tigris Imazethapyr 2SL** controls weeds by uptake by weed roots and translocation to the growing points where it stops weed growth. Susceptible weeds may emerge, growth will stop and the weeds will either die or are not competitive with the crop.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## SOIL APPLICATIONS WITH LIQUID FERTILIZERS

**Tigris Imazethapyr 2SL** can be applied to the soil in liquid fertilizers, alone or in combination with pendimethalin, or dimethenamide-P to soybeans. Mixtures including trifluralin may be applied to soybeans only. Follow all **Tigris Imazethapyr 2SL** label instructions regarding incorporation, timing of application, special instructions, restrictions and precautions. Apply treatments in 20 or more gals. of liquid fertilizer per acre with ground equipment. Always test the compatibility of **Tigris Imazethapyr 2SL** with the liquid fertilizer before mixing in the spray tank.

## PRE-EMERGENCE (SURFACE APPLICATIONS)

**Tigris Imazethapyr 2SL** offers flexibility in that it can be utilized in all production tillage systems. It can be applied prior to planting (up to 45 days prior to planting); at planting in conventional, reduced tillage or no-till production systems; or after planting and before crop emergence.

## NO-TILL OR REDUCED TILLAGE

Apply **Tigris Imazethapyr 2SL** treatments before, during or after planting. To ensure thorough coverage, use a minimum of 20 gals. of water per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues.

For maximum grass control, tank-mix **Tigris Imazethapyr 2SL** with pendimethalin, or dimethenamide-P. To kill existing vegetation, paraquat dichloride, glyphosate or 2,4-D (early pre-plant – see 2,4-D label for limitations) may be tank-mixed with **Tigris Imazethapyr 2SL** alone or in combination with pendimethalin, or dimethenamide-P. Delete paraquat dichloride, glyphosate or 2,4-D from the tank-mixture if vegetation is absent at the time of application.

**NOTE:** Adjust planters to ensure adequate soil coverage of seed.

## PRE-PLANT INCORPORATED APPLICATIONS

**Tigris Imazethapyr 2SL** may be applied following land preparation and must be thoroughly incorporated to a depth of 1 to 2 inches. If crops are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or a rolling cultivator. Maintain **Tigris Imazethapyr 2SL** in the surface 1 to 2 inches of the finished beds. Make application up to 45 days prior to planting soybeans.

When **Tigris Imazethapyr 2SL** is soil applied for control of nutsedge in peanuts, incorporate with two passes of the incorporation implement. Make the second pass at an offset angle to the first pass to minimize the potential for streaking.

## FEDERAL CONSERVATION RESERVE PROGRAM AND AGRICULTURAL RESERVE PROGRAM LAND SEEDED TO FORAGE LEGUME SPECIES AND PERENNIAL FORAGE GRASSES

### DIRECTIONS FOR USE

**Tigris Imazethapyr 2SL** is effective in controlling many annual broadleaf and grass weeds in CONSERVATION RESERVE PROGRAMS and AGRICULTURAL RESERVE PROGRAMS (SET-ASIDE) land seeded to forage legume or grass crops. An **Tigris Imazethapyr 2SL** application may result in temporary reduction in growth of legumes and grasses. Plants overcome temporary effects and become well established due to reduced weed competition.

### RESTRICTIONS

- DO NOT apply more than 4 fl. oz. **Tigris Imazethapyr 2SL** (0.0625 lb. ai) per application.
- DO NOT apply more than 4 fl. oz. **Tigris Imazethapyr 2SL** (0.0625 lb. ai) per acre per year.
- DO NOT make more than one application of **Tigris Imazethapyr 2SL** per year.
- DO NOT feed or graze legumes or grasses following an **Tigris Imazethapyr 2SL** application.
- DO NOT cut treated legumes or grasses for hay or forage.
- DO NOT harvest legume seed for livestock feed.
- DO NOT use seed from treated legumes for sprouting.
- DO NOT apply to seeded grasses until they have 4 leaves.

### COVER CROPS\*

#### LEGUMES:

Apply to forage legumes including alfalfa, Birdsfoot trefoil, clover, crown vetch, kudzu, lespedeza, lupin, milk vetch, sainfoin, trefoil, velvet bean, and vetch.

#### GRASSES:

**Tigris Imazethapyr 2SL** can be applied to the following grasses: big bluestem, little bluestem, switchgrass, Russian wild rye, intermediate wheatgrass, crested wheatgrass, western wheatgrass, tall wheatgrass, smooth brome, canarygrass, or orchardgrass.

\***NOTE:** Cover crops may be planted into fields previously treated with **Tigris Imazethapyr 2SL** for weed control in soybeans. In this case, do not make an **Tigris Imazethapyr 2SL** application to the cover crop until the following spring.

### POST-EMERGENCE APPLICATIONS OF TIGRIS IMAZETHAPYR 2SL TO CRP COVER CROPS

**APPLICATION RATE:** Apply **Tigris Imazethapyr 2SL** at 4 fl. oz./Acre (0.0625 lb. ai/Acre).

**APPLICATION TIMING:** **Tigris Imazethapyr 2SL** may be applied post-emergence to seedling legumes (with at least 3 fully expanded trifoliate leaves) or to established legumes. On established legumes, **Tigris Imazethapyr 2SL** may be applied in the fall or in the spring before weeds exceed the maximum specified size for control.

Refer to the **WEEDS CONTROLLED** under the **SOYBEAN** section of this label.



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## ALFALFA AND CLOVER

### DIRECTIONS FOR USE

**USE RATE:** 3 to 6 fl. oz./Acre (0.047 to 0.094 lb. ai/Acre)

Apply **Tigris Imazethapyr 2SL** at a broadcast rate of 3 to 6 fl. oz./Acre (0.047 to 0.094 lb. ai/Acre) post-emergence only to seedling or established alfalfa or clover grown for forage, hay, or seed.

### RESTRICTIONS

- DO NOT apply more than 6 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.094 lb. ai/Acre) per year to alfalfa or clover.
- DO NOT make more than one application of **Tigris Imazethapyr 2SL** per year.
- DO NOT apply **Tigris Imazethapyr 2SL** at more than 4 fl. oz./Acre (0.063 lb. ai/Acre) in North Dakota or Minnesota north of highway #210.
- DO NOT apply more than 4 fl. oz./Acre (0.063 lb. ai/Acre) to alfalfa or clover during the last year of the stand.
- DO NOT graze or harvest alfalfa or clover for 30 days following an application of **Tigris Imazethapyr 2SL** to alfalfa or clover.

### SEEDLING ALFALFA/CLOVER

**Tigris Imazethapyr 2SL** must be applied post-emergence to seedling alfalfa or clover. Apply **Tigris Imazethapyr 2SL** when the seedling alfalfa or clover is in the second (2nd) trifoliolate stage or larger and when the majority of the weeds are 1 to 3 inches. For low growing weeds (such as mustards) apply **Tigris Imazethapyr 2SL** before the rosette exceeds 3 inches. When **Tigris Imazethapyr 2SL** is applied to seedling alfalfa or clover, there may be a temporary reduction in growth.

### ESTABLISHED ALFALFA/CLOVER

**Tigris Imazethapyr 2SL** can be applied to established alfalfa or clover in the fall, in the spring to dormant, or semi-dormant alfalfa or clover (less than 3 inches of re-growth), or between cuttings. Any application must be made before significant alfalfa or clover growth or re-growth (3 inches) to allow **Tigris Imazethapyr 2SL** to reach the target weeds.

**Replanting:** If replanting is necessary in a field previously treated with **Tigris Imazethapyr 2SL**, do not plant alfalfa or clover for 4 months following a **Tigris Imazethapyr 2SL** application. Refer to the **ROTATIONAL CROP GUIDELINE** section on this label for plant-back interval of various crops.

### WEEDS CONTROLLED

When applied as directed, **Tigris Imazethapyr 2SL** will control or reduce competition from the weeds listed below. Refer to the **MIXING INSTRUCTIONS** section for specifications when weeds are at the maximum specified growth stage, or are under stress.

**NOTE:** R = Reduced Competition

Weeds noted with an "R" will be suppressed by **Tigris Imazethapyr 2SL**. For best results, apply before the weeds exceed the size indicated in the below table.

### BROADLEAF WEEDS CONTROLLED

Broadleaf Weeds	TIGRIS IMAZETHAPYR 2SL APPLICATION RATE		
	3 Fl. Oz./A (0.047 lb. ai/A)	4 Fl. Oz./A (0.063 lb. ai/A)	6 Fl. Oz./A (0.094 lb. ai/A)
	Maximum Weed Size (Inches)		
Artichoke, Jerusalem	R	6	8
Beets, wild	4	5	6
Bedstraw, catchweed	-	3	4
Buckwheat, wild	-	3	4
Chickweed, Common	R	3	4
Mouseear	R	3	3
Cocklebur, common	R	8	8
Cress, hoary	-	R	R
Dandelion	-	R	R(5)
Dock, broadleaf (seedling)	-	-	R(6)
curly (seedling)	-	-	R(6)
Dodder	-	-	R*
Fiddleneck	-	-	R(4)
Filaree, Redstem	-	R	3
Whitestem	-	R	3
Fleabane, rough	-	3	3



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557



**BROADLEAF WEEDS CONTROLLED (continued)**

Broadleaf Weeds	TIGRIS IMAZETHAPYR 2SL APPLICATION RATE		
	3 Fl. Oz./A (0.047 lb. ai/A)	4 Fl. Oz./A (0.063 lb. ai/A)	6 Fl. Oz./A (0.094 lb. ai/A)
	Maximum Weed Size (Inches)		
Flixweed	R	3	4
Goosefoot, nettleleaf	R	3	4
Groundsel, common	-	-	R(3)
Henbit	-	R	3
Jimsonweed	-	3	4
Knotweed, prostrate	-	R	3
Kochia (non-ALS resistant)	R	3	3
Lambsquarters, common (1 - 2 leaves)	-	R	R(2)
Lettuce, miner's	-	3	4
Mallow, Common	-	3	3
Little	-	3	3
Marshelder	-	4	6
Morningglory, Entireleaf	-	R	3
Ivyleaf	-	R	3
Pitted	-	R	3
Smallflower	R	3	4
Tall	-	R	3
Mustards, Tumble	3	3	4
Wild	3	3	4
Black	3	3	4
Nettle, burning	-	3	4
Nightshade, Black	3	3	4
Eastern black	3	3	4
Hairy	3	3	4
Oxtongue, bristly	-	-	R(3)
Pennycress, field	3	3	4
Pepperweed, field	3	3	4
Virginia	R	3	3
Pigweed, Redroot	4	6	8
Smooth	4	6	8
Spiny	-	6	8
Radish, wild	-	R	4
Ragweed, Common	-	2	3
Giant	-	3	3
Redmaids	-	3	4
Rocket, London	3	4	6
Yellow	R	3	4



**Manufactured for: Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**BROADLEAF WEEDS CONTROLLED (continued)**

Broadleaf Weeds	TIGRIS IMAZETHAPYR 2SL APPLICATION RATE		
	3 Fl. Oz./A (0.047 lb. ai/A)	4 Fl. Oz./A (0.063 lb. ai/A)	6 Fl. Oz./A (0.094 lb. ai/A)
	Maximum Weed Size (Inches)		
Rock purslane, desert	-	-	3
Shepherd's purse	3	3	4
Smartweed, Ladysthumb	R	3	4
Pennsylvania swamp (seedling)	R	3	4
	-	3	4
Spurge prostrate	-	R	3
spotted	-	R	3
petty	-	3	4
Spurry, corn	-	3	3
Sunflower, common	R	4	6
Swinecress	-	3	3
Tansymustard, green	3	3	4
pinnate	3	3	4
Thistle, Russian	R	3	3
Velvetleaf	R	3	4
Wartcress, creeping	-	2	3
Watercress	-	3	3
Willowweed, panicle	-	3	3

\*For best results in suppressing dodder (*Cuscuta* spp.), apply **Tigris Imazethapyr 2SL** with crop oil concentrate or methylated seed oil after dodder has emerged but prior to or soon after attachment.



**Manufactured for: Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**GRASSES AND SEDGES CONTROLLED**

Weeds Controlled*	TIGRIS IMAZETHAPYR 2SL APPLICATION RATE	
	4 Fl. Oz./A (0.063 lb. ai/A)	6 Fl. Oz./A (0.094 lb. ai/A)
	Maximum Weed Size (Inches)	
Barnyardgrass	R	3
Bluegrass, annual	–	R(3)
Canarygrass, littleseed	R	R(3)
Cereals, volunteer barley	R	R(4)
oats	R	R(4)
wheat	R	R(4)
Crabgrass, large	R	3
smooth	R	3
Cupgrass, woolly**	3	3
Foxtail, Giant	6	6
Green	3	4
Yellow	3	3
Johnsongrass, seedling	8	8
rhizome	R	R(6-12)
Millet, wild proso	R	3
Nutsedge, yellow	R	R(6)
purple	R	R(6)
Oats, wild	R	R(4)
Rice, red	3	4
Shattercane	8	10
Signalgrass, broadleaf	R	8
Quackgrass***	–	R(7)

\***Tigris Imazethapyr 2SL** is active against many grass species. However, when heavy grass pressure is anticipated, use **Tigris Imazethapyr 2SL** in a sequential application with a registered post-emergence grass herbicide such as sethoxydim for optimum control.

\*\***Tigris Imazethapyr 2SL** controls emerged woolly cupgrass only.

\*\*\*Quackgrass will be suppressed only when actively growing and before it exceeds 7 inches in height.

**TANK-MIX COMBINATIONS WITH OTHER HERBICIDES**

To control weeds not listed on the **Tigris Imazethapyr 2SL** label, herbicides such as bromoxynil, 2,4-DB, sethoxydim, rimsulfuron or clethodim may be tank-mixed with **Tigris Imazethapyr 2SL**. When **Tigris Imazethapyr 2SL** is used in combination with another herbicide, refer to the respective label for rates, methods of application, proper timing, weeds controlled, restrictions and precautions. Always use in accordance with the more restrictive label restrictions and precautions. DO NOT exceed label dosages.

**APPLICATION INFORMATION**

**Tigris Imazethapyr 2SL** is effective in controlling a broad spectrum of broadleaf and grass weeds. Alfalfa and clover are resistant to postemergence applications of **Tigris Imazethapyr 2SL** after the second trifoliate leaf has expanded. Minor height reduction or slight leaf yellowing may occur soon after application.

Apply **Tigris Imazethapyr 2SL** as an early post-emergence treatment when the weeds are actively growing. Weeds are generally easier to control before they exceed 3 inches in height. Weeds under stress are less susceptible to control in cold or drought stress conditions.

If applied to alfalfa or clover under cool conditions (40°F or less), temporary stunting and yellowing of the crop may occur.

**Stand Establishment**

Apply **Tigris Imazethapyr 2SL** after the alfalfa or clover has 2 fully expanded trifoliate leaves. Weeds must not exceed the size listed in the WEEDS CONTROLLED tables. **Tigris Imazethapyr 2SL** may be applied to summer, fall or spring seeded alfalfa or clover



**Manufactured for: Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

### Inter-seeded Oats

Oats inter-seeded with alfalfa will reduce soil erosion and allow the alfalfa or clover to establish. Oats, however can compete with the alfalfa or clover. An application of **Tigris Imazethapyr 2SL** will kill or significantly reduce the growth of the oats and allow the alfalfa or clover to establish with minimal erosion or competition from the oats. Apply **Tigris Imazethapyr 2SL** to the oats when they have 3 to 4 leaves.

### ESTABLISHED ALFALFA / CLOVER – DORMANT

Apply **Tigris Imazethapyr 2SL** to dormant alfalfa or clover in the fall following the last cutting. Apply **Tigris Imazethapyr 2SL** in the spring to dormant alfalfa or clover, or as alfalfa or clover breaks dormancy. Apply spring treatments prior to excessive alfalfa or clover growth (less than 3 inches), to reduce spray interference.

### ESTABLISHED ALFALFA / CLOVER – GROWING

For weed control during the season, apply **Tigris Imazethapyr 2SL** following alfalfa or clover cutting. Remove the hay from the field and apply **Tigris Imazethapyr 2SL** prior to excessive alfalfa or clover regrowth.

### Perennial Grass Suppression

If perennial grasses (such as orchardgrass, fescues, bromes or timothy) are present in an alfalfa or clover stand, **Tigris Imazethapyr 2SL** will reduce the growth and competitive effect of the grass.

### NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, LIMA, AND SMALL WHITE TYPE DRY BEANS, ADZUKI, LENTILS, WHITE LUPINS, CHICKPEAS (GARBANZO BEANS), DRY EDIBLE PEAS, ENGLISH AND SOUTHERN PEAS

**DIRECTIONS FOR USE** in the states east of and including: North Dakota, South Dakota, Wyoming, Colorado, and New Mexico (except the states east of and including: Vermont, Massachusetts, and Connecticut). Refer to map for geographical use area.



Use only non-ionic surfactants as a spray additive for post-emergence applications. **DO NOT** use crop oils, methylated seed oils, or petroleum oils.

### RESTRICTIONS

- DO NOT make more than one application of **Tigris Imazethapyr 2SL** per year.
- DO NOT apply more than 4 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.047 lb. ai/Acre) per application.
- DO NOT apply more than 4 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.063 lb. ai/Acre) per year to peas and beans in this region.
- Allow at least 30 days between application and harvest of succulent lima beans, English peas, and Southern peas. Allow at least 60 days between application and harvest of dry edible peas, lentils, chickpeas, and other dry bean or pea types listed on this label.
- DO NOT APPLY **Tigris Imazethapyr 2SL** POST-EMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR PEAS ARE AT LEAST THREE INCHES IN HEIGHT OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.
- DO NOT APPLY **Tigris Imazethapyr 2SL** POST-EMERGENCE TO LIMA BEANS, LENTILS, WHITE LUPINS, OR CHICKPEAS.
- DO NOT apply to Domino variety black turtle beans.
- DO NOT apply this product through any type of irrigation system.

Pinto varieties UI-111 and Olathe are more sensitive to **Tigris Imazethapyr 2SL** than other varieties.

### APPLICATION INSTRUCTIONS

#### NAVY, GREAT NORTHERN, RED KIDNEY, BLACK TURTLE, CRANBERRY, PINTO, AND SMALL WHITE DRY BEANS, ADZUKI, DRY EDIBLE PEAS, ENGLISH AND SOUTHERN PEAS

In Michigan or the Delaware, Maryland, and Virginia (Delmarva) peninsula: DO NOT apply more than 2 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.031 lb. ai/Acre) to sands or loamy sand soils.

In North Dakota or north of Highway 210 in Minnesota: DO NOT apply more than 2 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.031 lb. ai/Acre).

#### Pre-Plant Incorporated Applications:

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of up to 3 fl. oz./Acre (0.047 lb. ai) to dry beans (navy, great northern, red kidney, black turtle, cranberry, pinto and small white type dry beans, and adzuki), dry edible peas, and English peas, or up to 4 fl. oz./Acre (0.063 lb. ai/Acre) for southern peas only, within 1 week before planting. Applied pre-plant incorporated, **Tigris Imazethapyr 2SL** may be tank-mixed with a registered grass herbicide.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**Pre-Emergence Applications:**

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of up to 3 fl. oz./Acre (0.047 lb. ai/Acre) to dry beans, dry edible peas and English peas, or up to 4 fl. oz./Acre (0.063 lb. ai/Acre) for southern peas only, immediately after, or up to 3 days after planting. **Tigris Imazethapyr 2SL** may be applied in a tank-mix with a registered grass herbicide or applied pre-emergence following a pre-plant incorporated application of a registered grass herbicide.

**Early Post-Emergence Applications:**

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of up to 3 fl. oz./Acre (0.047 lb. ai/Acre) to dry beans, dry edible peas, and English peas, or up to 4 fl. oz./Acre (0.063 lb. ai/Acre) for southern peas only. Apply to dry beans with at least one fully expanded trifoliolate leaf. Apply to dry edible peas, English peas, and southern peas at least 3 inches in height but prior to 5 nodes and before flowering. The use of trifluralin prior to **Tigris Imazethapyr 2SL** application may increase the likelihood and severity of crop injury. A non-ionic surfactant must be added to the spray solution. The non-ionic surfactant must contain at least 80% active ingredient and shall be used at a rate of 2 pts. per 100 gals. of spray mixture.

Sodium bentazon may be tank-mixed with **Tigris Imazethapyr 2SL** to control weeds not listed on the **Tigris Imazethapyr 2SL** label. Addition of sodium bentazon may also cause antagonism, thereby reducing control of grass weeds. Nitrogen-based fertilizer may be included as a spray additive ONLY when **Tigris Imazethapyr 2SL** is tank-mixed with sodium bentazon. Refer to the sodium bentazon label for proper application rates and restrictions. Always use in accordance with the more restrictive label restrictions and precautions.

**LIMA BEANS, CHICKPEAS (GARBANZOS), LENTILS, AND WHITE LUPINS**

**DO NOT** apply **Tigris Imazethapyr 2SL** to white lupins grown on sand or loamy sand soils.

In Michigan or the Delaware, Maryland, and Virginia (Delmarva) peninsula: **DO NOT** apply more than 2 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.031 lb. ai/Acre) to sands or loamy sand soils.

In North Dakota or north of Highway 210 in Minnesota: **DO NOT** apply more than 2 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.031 lb. ai/Acre).

**Pre-Plant Incorporated Applications:**

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of up to 3 fl. oz./Acre (0.047 lb. ai/Acre) within 1 week before planting. Applied pre-plant incorporated, **Tigris Imazethapyr 2SL** may be tank-mixed with a registered grass herbicide.

**Pre-Emergence Applications:**

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of up to 3 fl. oz./Acre (0.047 lb. ai/Acre) immediately after or up to 3 days after planting. **Tigris Imazethapyr 2SL** may be applied in a tank-mix with a registered grass herbicide or applied pre-emergence following a pre-plant incorporated application of a registered grass herbicide.

**WEEDS CONTROLLED**

**Tigris Imazethapyr 2SL** applied at the broadcast rate of 2 fl. oz./Acre (0.031 lb. ai/Acre) pre-plant incorporated, pre-emergence, or early post-emergence will control:

Mustard, wild

Nightshade, black\*

Nightshade, Eastern black\*

\*suppression only

**Tigris Imazethapyr 2SL** applied at the broadcast rate of 3 fl. oz./Acre (0.047 lb. ai/Acre) pre-plant incorporated, pre-emergence, or early post-emergence will control:

Mustard, wild

Nightshade, black

Nightshade, Eastern black

Nightshade, hairy

Pigweed, redroot

Post-emergence applications of 3 fl. oz./Acre (0.047 lb. ai/Acre) must be made to weeds less than 2 inches tall for best results.



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

When applied as directed at the broadcast rate of 4 fl. oz./Acre (0.063 lb. ai/Acre) for southern peas only, **Tigris Imazethapyr 2SL** will control or reduce competition from the weeds listed below:

**NOTE:** C = Control, R = Reduced Competition

The number under **Maximum Leaf Stage** indicates the MAXIMUM number of leaves at which weeds must be sprayed post-emergence.

**BROADLEAF WEEDS CONTROLLED**

Broadleaf Weeds	SOIL APPLIED	POST-EMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem	-	8	6-10
Bristly starbur	-	2	1-2
Buffalobur	C*	-	-
Carpetweed	C	-	-
Cocklebur, common	C*	8	1-8
Galinsoga	C		
Jimsonweed	C**	4	1-3
Kochia (non-ALS resistant)	C	4	1-3
Lambsquarters, common	C**	R	1-2
Mallow, Venice	R	2	1-2
Morningglory, Entireleaf	R	2	1-2
Ivyleaf	R	2	1-2
Pitted	R	2	1-2
Smallflower	C	4	1-3
Tall	R	2	1-2
Mustard sp.	C	4	1-3
Nightshade, Black	C	4	1-3
Eastern black	C	4	1-3
Hairy	C	4	1-3
Pigweed, Redroot	C	4	1-4
Smooth	C	4	1-4
Spiny	C	4	1-4
Poinsettia, wild	C	-	-
Puncturevine	C	-	-
Purslane, common	C	-	-
Pusley, Florida	C	-	-
Ragweed, Common	R	4	1-3
Giant	R	4	1-3
Sage, barnyard	-	R	1-3
Sida, prickly	C**	-	-
Smartweed, Ladysthumb	C	4	1-3
Pennsylvania	C	4	1-3
Spurge prostrate	C	4	1-3
spotted	C	4	1-3



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**BROADLEAF WEEDS CONTROLLED (continued)**

Broadleaf Weeds	SOIL APPLIED	POST-EMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Sunflower, common	C**	4	1-3
Thistle, Canada	-	R	1-3
Velvetleaf	C**	3	1-3

\*Use soil applications for light to moderate infestations only. Must be pre-plant incorporated for best results.

\*\*When soil applied, common lambsquarters, jimsonweed, prickly sida, velvetleaf, and common sunflower are more consistently controlled by preplant incorporated treatments.

**GRASS WEEDS CONTROLLED\***

Broadleaf Weeds	SOIL APPLIED	POST-EMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Barnyardgrass	R	3	1-3
Crabgrass, large smooth	R	3	1-3
	R	3	1-3
Cupgrass, woolly	-	3**	1-3
Foxtail, giant green robust purple robust white yellow	C	6	1-6
	C	3	1-3
	C	3	1-3
	C	3	1-3
	C	3	1-3
Goosegrass	R	-	-
Johnsongrass, seedling rhizome	C	6	1-8
	-	R	1-8
Panicum, fall Texas	R	-	-
	R	-	-
Red rice	-	3	-
Shattercane	R	6	-
Signalgrass, broadleaf	R	4	1-8

\*When soil applied to grasses, more consistent control can be obtained from pre-plant incorporated treatments.

\*\*Tigris Imazethapyr 2SL controls emerged woolly cupgrass only.

**SEDGES CONTROLLED**

Grass Weeds	SOIL APPLIED	POST-EMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Nutsedge, purple yellow	R	R	1-3
	R	R	-

**DO NOT** count cotyledon leaves when determining weed stage of growth.

Refer to the **ADDITIONAL USE INFORMATION** section for additional instructions.



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## RED KIDNEY BEANS

**DIRECTIONS FOR USE** in the state of California.

### APPLICATION RATE AND TIMING

#### Post-Emergence Applications:

Apply **Tigris Imazethapyr 2SL** at the rate of 3 fl. oz./Acre (0.047 lb. ai/Acre). A non-ionic surfactant must be added to the spray solution. The non-ionic surfactant must contain at least 80% active ingredient and used at a rate of 2 pts. per 100 gals. of spray mixture.

Apply **Tigris Imazethapyr 2SL** herbicide when weeds are actively growing and red kidney beans have at least 1 fully expanded trifoliolate leaf.

For maximum weed control, cultivate 7 – 10 days following a post-emergence **Tigris Imazethapyr 2SL** application. This timely cultivation will enhance residual weed control, especially under dry conditions.

### RESTRICTIONS

- DO NOT MAKE MORE THAN ONE APPLICATION OF **Tigris Imazethapyr 2SL** PER YEAR.
- DO NOT apply more than 3 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.047 lb. ai/Acre) per application.
- DO NOT apply more than 3 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.047 lb. ai/Acre) per year to Red Kidney Beans.
- DO NOT APPLY **Tigris Imazethapyr 2SL** POST-EMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.
- DO NOT apply **Tigris Imazethapyr 2SL** post-emergence when the crop and weeds have been subjected to stress conditions such as temperature or moisture extremes.
- DO NOT apply by aerial application.
- Allow at least 60 days between application and harvest.

### WEEDS CONTROLLED

When applied as directed, **Tigris Imazethapyr 2SL** will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section for instructions when weeds are at the maximum specified growth stage, or are under stress. (The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds need to be sprayed post-emergence.)

Weeds Controlled	POST-EMERGENCE	
	Maximum Leaf Stage	Size (Inches)
Kochia (non-ALS resistant)	4	1-3
Mustard sp.	4	1-3
Nightshade, black	4	1-3
Eastern black hairy	4	1-3
Pigweed, redroot	4	1-2
		1-3

Refer to the **ADDITIONAL USE INFORMATION** section for additional instructions.

## SNAP BEANS

**DIRECTIONS FOR USE** in the states of Alabama, Florida, Georgia, Illinois, Minnesota, Michigan, New Jersey, North Carolina, and Wisconsin.

### RESTRICTIONS

- DO NOT apply by aerial application.
- DO NOT apply **Tigris Imazethapyr 2SL** after July 31st (June 20th in New Jersey).
- DO NOT MAKE MORE THAN ONE APPLICATION OF **Tigris Imazethapyr 2SL** PER YEAR.
- DO NOT apply more than 1.5 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.023 lb. ai/Acre) per application.
- DO NOT apply more than 1.5 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.023 lb. ai/Acre) per year to Snap Beans.
- Allow at least 30 days between application and harvest.

### APPLICATION INSTRUCTIONS

#### Pre-Plant Incorporated Applications:

Apply **Tigris Imazethapyr 2SL** at 1.5 fl. oz./Acre (0.023 lb. ai/Acre) within 1 week of planting. Applied pre-plant incorporated, **Tigris Imazethapyr 2SL** may be tank-mixed with a registered grass herbicide.

#### Pre-Emergence Applications:

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of 1.5 fl. oz./Acre (0.023 lb. ai/Acre) immediately after, or up to 1 day after planting. **Tigris Imazethapyr 2SL** may be applied in a tank-mix with a registered grass herbicide or applied pre-emergence following a pre-plant incorporated application of a registered grass herbicide.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557



## WEEDS SUPPRESSED

**Tigris Imazethapyr 2SL** applied at the broadcast rate of 1.5 fl. oz./Acre (0.023 lb. ai/Acre) pre-plant incorporated or pre-emergence will suppress or reduce competition of the following weeds:

Common Purslane  
Eastern Black Nightshade  
Redroot Pigweed  
Wild Mustard

Refer to the **ADDITIONAL USE INFORMATION** section for additional instructions.

## SNAP BEANS

**DIRECTIONS FOR USE** in the states of Arkansas, Missouri, North Carolina, Oklahoma, Texas (counties of Bailey, Castro, Lamb and Parmer only), and New Mexico (counties of Curry and Roosevelt only).

### RESTRICTIONS

- DO NOT apply by aerial application.
- DO NOT apply **Tigris Imazethapyr 2SL** after July 31st.
- DO NOT MAKE MORE THAN ONE APPLICATION OF **Tigris Imazethapyr 2SL** PER YEAR.
- DO NOT apply more than 1.5 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.023 lb. ai/Acre) per application.
- DO NOT apply more than 1.5 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.023 lb. ai/Acre) per year to Snap Beans.
- Allow at least 30 days between application and harvest.
- DO NOT APPLY **Tigris Imazethapyr 2SL** POST-EMERGENCE BEFORE CROP HAS AT LEAST ONE TRUE LEAF OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.

### APPLICATION INSTRUCTIONS

#### Post-Emergence Applications:

Apply **Tigris Imazethapyr 2SL** at 1.5 fl. oz./Acre (0.023 lb. ai/Acre) in a tank-mix combination with sodium bentazon. A non-ionic surfactant must be added to the spray solution. The non-ionic surfactant must contain at least 80% active ingredient and used at a rate of 2 pts. per 100 gals. of spray mixture.

Refer to the sodium bentazon label for proper application rates and restrictions.

## WEEDS SUPPRESSED

**Tigris Imazethapyr 2SL** applied at the broadcast rate of 1.5 fl. oz./Acre (0.023 lb. ai/Acre) post-emergence will suppress or reduce competition of the following weeds:

Eastern Black Nightshade  
Redroot Pigweed

Refer to the **ADDITIONAL USE INFORMATION** section for additional instructions.

## SUCCULENT PEAS, DRY EDIBLE PEAS, LENTILS, CHICKPEAS, AND LIMA BEANS

**DIRECTIONS FOR USE** in the states of Idaho, Montana, Nevada, Oregon, Utah, and Washington.

### APPLICATION RATE AND TIMING

#### Pre-Plant Applications for No-Till and Minimum Tillage Systems Only:

Apply **Tigris Imazethapyr 2SL** at a broadcast rate of 3 fl. oz./Acre (0.047 lb. ai/Acre) within 30 days before planting. If incorporated, DO NOT incorporate deeper than 3 inches.

In no-till and minimum tillage systems, apply **Tigris Imazethapyr 2SL** in the fall prior to spring planting. Rainfall is required for incorporation and activation. Unpredictable weed control can be expected since factors such as length of time between application and planting as well as uncontrollable weather factors will determine herbicide activity and longevity. Apply **Tigris Imazethapyr 2SL** in the fall when soil temperature at the 4-inch depth is less than 55°F and before the ground is frozen.

#### Pre-Plant Incorporated Applications:

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of 3 fl. oz./Acre (0.047 lb. ai/Acre) within 1 week before planting. DO NOT incorporate deeper than 3 inches.

#### Pre-Emergence Applications:

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of 3 fl. oz./Acre (0.047 lb. ai/Acre) after planting, but prior to crop emergence.

Tank mix **Tigris Imazethapyr 2SL** with metribuzin to assist in the control of lambsquarters or mayweed-chamomile (dogfennel). Refer to the metribuzin label for proper application rates and restrictions.

#### Post-Emergence Applications (Dry Edible Peas Only):

Apply **Tigris Imazethapyr 2SL** at 2 fl. oz./Acre (0.031 lb. ai/Acre). A non-ionic surfactant must be added to the spray solution. The nonionic surfactant must contain at least 80% active ingredient and used at a rate of 2 pts. per 100 gals. of spray mixture.

Sodium bentazon may be tank-mixed with **Tigris Imazethapyr 2SL** to control weeds not listed on the **Tigris Imazethapyr 2SL** label. Addition of sodium bentazon may also cause antagonism, thereby reducing control of grass weeds. Nitrogen-based fertilizer may be included as a spray additive only when **Tigris Imazethapyr 2SL** is tank-mixed with sodium bentazon. Use liquid fertilizer at 1.25 to 2.5 gals. per 100 gals. of spray solution or ammonium sulfate at the rate of 12 – 15 lbs./100 gals. of spray solution.



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## RESTRICTIONS

- DO NOT apply more than 3 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.047 lb. ai/Acre) per application to Peas and Beans in this region.
- DO NOT apply more than 3 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.047 lb. ai/Acre) per year to Peas and Beans in this region.
- DO NOT MAKE MORE THAN ONE APPLICATION OF **Tigris Imazethapyr 2SL** PER YEAR.
- DO NOT APPLY **Tigris Imazethapyr 2SL** POST-EMERGENCE BEFORE CROP HAS AT LEAST ONE TRIFOLIATE LEAF OR PEAS ARE AT LEAST THREE INCHES IN HEIGHT OR CROP INJURY (REDUCED CROP GROWTH AND/OR DELAYED MATURITY) MAY RESULT.
- DO NOT APPLY **Tigris Imazethapyr 2SL** POST-EMERGENCE TO LIMA BEANS, LENTILS, OR CHICKPEAS.
- Allow at least 30 days between application and harvest for succulent peas and succulent lima beans.
- Allow at least 60 days between application and harvest for dry edible peas, chickpeas, lentils, and dry lima beans.

## WEEDS CONTROLLED

**Tigris Imazethapyr 2SL** applied PPI and/or Pre-emergence at 3 fl. oz./Acre (0.047 lb. ai/Acre) will control:

Weeds Controlled	Pre-Plant Incorporated	Pre-Emergence
Buckwheat, wild	C	C
Kochia (non-ALS resistant)	C	C
Lambsquarters, common	C	-
Mustard, wild	C	C
Nightshade, black	C	C
Eastern black	C	C
hairy	C	C
Pigweed, redroot	C	C
Shepherd's purse	C	C
Thistle, Russian	C	C

**NOTE:** C = Control

**Tigris Imazethapyr 2SL** applied post-emergence at the broadcast rate of 2 fl. oz./Acre (0.031 lb. ai/Acre) will control:

Black nightshade\*

Eastern black nightshade\*

Hairy nightshade\*

Wild mustard

\*Suppression only

Refer to the **ADDITIONAL USE INFORMATION** section for additional instructions.

## CHICKPEAS

**DIRECTIONS FOR USE** in the states of Arizona and California.

### APPLICATION RATE AND TIMING

#### Pre-Plant Incorporated Applications:

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of up to 3 fl. oz./Acre (0.047 lb. ai/Acre) within 1 week before planting. Applied pre-plant incorporated, **Tigris Imazethapyr 2SL** may be tank-mixed with a registered grass herbicide.

#### Pre-Emergence Applications:

Apply **Tigris Imazethapyr 2SL** at the broadcast rate of up to 3 fl. oz./Acre (0.047 lb. ai/Acre) immediately after or up to 3 days after planting. **Tigris Imazethapyr 2SL** may be applied in a tank-mix with a registered grass herbicide or applied pre-emergence following a pre-plant incorporated application of a registered grass herbicide.

## RESTRICTIONS

- DO NOT apply more than 3 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.047 lb. ai/Acre) per application to Chickpeas in this region.
- DO NOT apply more than 3 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.047 lb. ai/Acre) per year to Chickpeas in this region.
- DO NOT MAKE MORE THAN ONE APPLICATION OF **Tigris Imazethapyr 2SL** PER YEAR.
- Allow at least 30 days between application and harvest of succulent chickpeas. Allow at least 60 days between application and harvest of dry chickpeas.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## WEEDS CONTROLLED

Weeds Controlled	Pre-Plant Incorporated	Pre-Emergence
Buckwheat, wild	C	C
Kochia (non-ALS resistant)	C	C
Lambsquarters, common	C	-
Mustard, wild	C	C
Nightshade, black	C	C
Eastern black	C	C
hairy	C	C
Pigweed, redroot	C	C
Shepherd's purse	C	C
Thistle, Russian	C	C

**NOTE:** C = Control

Refer to the **ADDITIONAL USE INFORMATION** section for additional instructions.

## PEANUTS

**(Not for use in California)**

### DIRECTIONS FOR USE

**USE RATE:** 4 fl. oz./Acre (0.063 lb. ai/Acre)

Apply **Tigris Imazethapyr 2SL** at a broadcast rate of 4 fl. oz./Acre (0.063 lb. ai/Acre) for all methods of application (except sequential – see below): pre-plant incorporated, pre-emergence, ground-cracking and post-emergence. At this broadcast rate, 1 gal. of **Tigris Imazethapyr 2SL** will treat 32 acres of peanuts.

**Tigris Imazethapyr 2SL** may also be applied in a sequential application. Apply 2 fl. oz./Acre (0.031 lb. ai/Acre) in a soil application (pre-plant incorporated or preemergence) followed by 2 fl. oz./Acre (0.031 lb. ai/Acre) applied at ground-crack or post-emergence.

### RESTRICTIONS

- DO NOT apply more than 4 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.063 lb. ai/Acre) per application.
- DO NOT apply more than 4 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.063 lb. ai/Acre) per year to Peanuts.
- DO NOT make more than one application per year at the 4 fl. oz./Acre rate (0.063 lb. ai/Acre) or two applications at the 2 fl. oz./Acre rate (0.031 lb. ai/Acre).
- DO NOT graze or feed treated peanut forage, vines, hay or straw to livestock.
- There must be an interval of at least 85 days between an application of **Tigris Imazethapyr 2SL** and peanut harvest.
- DO NOT apply pendimethalin + imazethapyr herbicides to peanuts the same year as **Tigris Imazethapyr 2SL**.

**NOTE:** In Arizona for use only in Yuma and La Paz counties.



**Manufactured for: Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**WEEDS CONTROLLED**

When applied as directed, **Tigris Imazethapyr 2SL** will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section for instructions when weeds are at the maximum specified growth stage, or are under stress. (The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds are to be sprayed post-emergence.)

**NOTE:** C = Control, R = Reduced Competition

**BROADLEAF WEEDS CONTROLLED**

Broadleaf Weeds	SOIL APPLIED	AT-CRACK	POST-EMERGENCE	
			Maximum Leaf Stage	Size (Inches)
Spurge prostrate	C	C	4	1-3
spotted	C	C	4	
toothed	C	C	-	-
Sunflower	C*	C	4	1-3
Velvetleaf	C*	C	4	1-3

\*When **Tigris Imazethapyr 2SL** is soil applied, these weeds are more consistently controlled by pre-plant incorporated treatments.

**GRASS WEEDS CONTROLLED**

Broadleaf Weeds	SOIL APPLIED*	AT-CRACK	POST-EMERGENCE	
			Maximum Leaf Stage	Size (Inches)
Barnyardgrass	R	R	3	1-3
Crabgrass, large	R	C	3	1-3
smooth	R	C	3	1-3
Cupgrass, woolly	-	-	3	1-3
Foxtail, giant	C		6	1-6
green	C		3	1-3
yellow	C		3	1-3
Goosegrass	R		-	-
Johnsongrass, seedling	C	C	6	1-8
rhizome	-	-	R	6-12
Panicum, fall	R	-	-	-
Texas	R	-	-	-
Red rice	-	-	3	1-3
Shattercane	R	R	6	1-8
Signalgrass, broadleaf	R	C	4	1-6

\*When **Tigris Imazethapyr 2SL** is soil applied to grasses, more consistent control can be obtained from pre-plant incorporated treatments.

**SEDGES**

Weeds Controlled	SOIL APPLIED*	AT-CRACK	POST-EMERGENCE	
			Maximum Leaf Stage	Size (Inches)
Nutsedge, purple	C	C	3	1-3
yellow	C	C	3	1-3

\*When **Tigris Imazethapyr 2SL** is soil applied to grasses, more consistent control can be obtained from pre-plant incorporated treatments.

**DO NOT** count cotyledon leaves when determining weed stage of growth.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**BROADLEAF WEEDS CONTROLLED (continued)**

Broadleaf Weeds	SOIL APPLIED	AT-CRACK	POST-EMERGENCE	
			Maximum Leaf Stage	Size (Inches)
Alligator weed	-	C	4	1-3
Anoda, spurred	C	C	2	1-2
Bristly starbur	-	-	2	1-2
Buffalobur	C*	C	R	1-3
Carpetweed	C	C	-	-
Cocklebur, common	R	C	8	1-8
Devil's claw	C	C	-	-
Galinsoga	C	C	-	-
Jimsonweed	C*	C	4	1-3
Lambsquarters, common	C*	C	R	1-2
Morningglory, Entireleaf	R	C	2	1-2
Ivyleaf	R	C	2	1-2
Pitted	R	C	2	1-2
Smallflower	C	C	4	1-3
Tall	R	C	2	1-2
Mustard sp.	C	C	4	1-3
Nightshade, black	C	C	4	1-3
Eastern black hairy	C	C	4	1-3
Pigweed, redroot	C	C	8	1-8
smooth	C	C	8	1-8
spiny	C	C	8	1-8
Poinsettia, wild	C	C	-	-
Puncturevine	C	C	-	-
Purslane, common	C	C	-	-
Pusley, Florida	C	C	-	-
Ragweed, common	R	R	4	1-3
giant	R	R	4	1-3
Sida, prickly (teaweed)	C*	C	-	-
Smartweed, Ladysthumb	C	C	4	1-3
Pennsylvania	C	C	4	1-3


**Manufactured for: Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**"AT-CRACK"** APPLICATION refers to the time when the soil cracks due to the emerging peanut seedling. This generally occurs from 10 to 14 days following planting. At this time weeds have generally not germinated, or are in the seedling stage. If weeds have more than 2 true leaves, refer to the **POST-EMERGENCE** weed control column for weeds controlled.

In West Texas and New Mexico, wait until late cracking (most of the peanuts have emerged) before applying **Tigris Imazethapyr 2SL**.

**Tigris Imazethapyr 2SL** is active against many broadleaf and grass species. However, when heavy grass or common lambsquarters pressure is anticipated, apply **Tigris Imazethapyr 2SL** in combination with a registered soil-applied grass herbicide (See HERBICIDE COMBINATIONS section).

#### WEEDS CONTROLLED BY SEQUENTIAL APPLICATIONS OF TIGRIS IMAZETHAPYR 2SL

The sequential (split) application of **Tigris Imazethapyr 2SL** consists of an application of 2 fl. oz./Acre (0.031 lb. ai/Acre) of product soil applied (either pre-plant incorporated or pre-emergence) followed by 2 fl. oz./Acre (0.031 lb. ai/Acre) applied either at ground-crack or post-emergence.

When applied as a sequential treatment, **Tigris Imazethapyr 2SL** will control the weeds listed under the "SOIL APPLIED" and "AT-CRACK" applications in the **BROADLEAF WEEDS** and **GRASS WEEDS** tables (in the **PEANUTS** section of the label). It enhances the control of yellow and purple nutsedge. Apply the second application before the nutsedge exceeds 3 leaves.

#### HERBICIDE COMBINATIONS

##### GRASS WEEDS

When applied as directed, **Tigris Imazethapyr 2SL** pre-plant incorporated or pre-emergence combination treatments with pendimethalin, trifluralin, alachlor, S-metolachlor, isoxaflutole, or ethalfluralin will control the weeds listed in following table, in addition to those controlled by **Tigris Imazethapyr 2SL** alone.

GRASSES	Pendimethalin <sup>a</sup>	Trifluralin <sup>b</sup>	Alachlor	S-metolachlor	Isoxaflutole <sup>b</sup>	Ethalfluralin <sup>b</sup>
Barnyardgrass	X	X	X	X	X	X
Crabgrass, smooth	X	X	X	X	X	X
Crabgrass, large	X	X	X	X	X	X
Crowfootgrass	X	X	-	-	X	-
Goosegrass	X	X	X	X	X	X
Panicum, fall	X	X	X	X	X	X
Panicum, Texas	X	X	-	-	X	X
Sandbur, field	X	X	-	-	X	X
Signalgrass, broadleaf	X <sup>b</sup>	X	X	X	X	X
Witchgrass	X	X	X	X	-	X

<sup>a</sup> Pre-plant incorporated tank-mixture applications of **Tigris Imazethapyr 2SL** plus pendimethalin will suppress the growth of itchgrass and rhizome johnsongrass.

<sup>b</sup> Pre-plant incorporated treatments only.

A selective post-emergence grass herbicide such as sethoxydim or fenoxaprop-p-ethyl may be mixed with **Tigris Imazethapyr 2SL** to control grasses not controlled by **Tigris Imazethapyr 2SL**. In some cases the activity of the grass herbicide may be reduced when mixed with **Tigris Imazethapyr 2SL**. The reduction in activity may be overcome by delaying the application of the post-emergence grass herbicide 7 days following the application of **Tigris Imazethapyr 2SL**. If the post-emergence grass herbicide is applied first, wait 3 days before applying **Tigris Imazethapyr 2SL**. Refer to the respective grass herbicide label for directed application rate, weed size and restrictions.

#### BROADLEAF WEEDS

Broadleaf herbicides that can be tank-mixed with **Tigris Imazethapyr 2SL** include sodium bentazon, acifluorfen, flufenacet and 2,4-DB.

For the control of sicklepod, morningglories, prickly sida and common ragweed, add 2,4-DB to the **Tigris Imazethapyr 2SL** spray mixture. For the control of Florida beggarweed, add flufenacet to the spray mixture. Refer to the 2,4-DB or flufenacet label for specific directions for use, application rates and restrictions.

**Tigris Imazethapyr 2SL** may also be applied post-emergence in tank-mixture with chlorothalonil, acephate, or boron.

#### SOYBEANS

(Not for use in California)

#### DIRECTIONS FOR USE

**USE RATE:** 4 fl. oz./Acre (0.063 lb. ai/Acre)

Apply **Tigris Imazethapyr 2SL** at a broadcast rate of 4 fl. oz./Acre (0.063 lb. ai/Acre) for all methods of application: early pre-plant, pre-plant incorporated, pre-emergence, and post-emergence (including minimum and no-till). At this broadcast rate, 1 gal. of **Tigris Imazethapyr 2SL** will treat 32 acres of soybeans. (See instructions under section **APPLICATIONS TO SOYBEANS IN NORTH DAKOTA AND MINNESOTA** for applications in North Dakota and Minnesota north of highway #210.). Make **Tigris Imazethapyr 2SL** applications before soybean bloom.



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**RESTRICTIONS**

- DO NOT apply more than 4 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.063 lb. ai/Acre) per application.
- DO NOT apply more than 4 fl. oz./Acre of **Tigris Imazethapyr 2SL** (0.063 lb. ai/Acre) per year to Soybeans.
- DO NOT make more than one application per year.
- There must be an interval of at least 85 days between an application of **Tigris Imazethapyr 2SL** and soybean harvest.
- DO NOT graze or feed treated soybean forage, hay or straw to livestock.
- DO NOT tank-mix **Tigris Imazethapyr 2SL** with clomazone containing herbicides. **Tigris Imazethapyr 2SL** may be applied post-emergence following a soil application of a clomazone containing herbicide.

**WEEDS CONTROLLED**

When applied as directed, **Tigris Imazethapyr 2SL** will control or reduce competition from the weeds listed below. Refer to the MIXING INSTRUCTIONS section for instructions when weeds are at the maximum specified growth stage, or are under stress. (The number under Maximum Leaf Stage indicates the MAXIMUM number of leaves at which weeds need to be sprayed post-emergence.)

**NOTE:** C = Control, R = Reduced Competition

**DO NOT** count cotyledon leaves when determining weed stage of growth.

**BROADLEAF WEEDS CONTROLLED**

Broadleaf Weeds	SOIL APPLIED	POST-EMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Alligator weed	-	4	1-3
Anoda, spurred	C	2	1-2
Artichoke, Jerusalem	-	8	6-10
Bristly starbur	-	2	1-2
Buffalobur	C*	R	1-3
Carpetweed	C	-	-
Cocklebur, common	R	8	1-8
Galinsoga	C	-	-
Jimsonweed	C*	4	1-3
Kochia (non-ALS resistant)	C	4	1-3
Lambsquarters, common	C*	R	1-2
Mallow, Venice	R	-	-
Marshelder	C	4	1-3
Morningglory, entireleaf	R	2	1-2
ivyleaf	R	2	1-2
pitted	R	2	1-2
smallflower	C	4	1-3
tall	R	2	1-2
Mustard sp.	C	4	1-3
Nightshade, black	C	4	1-3
Eastern black	C	4	1-3
hairy	C	4	1-3
Pigweed, redroot	C	8	1-8
smooth	C	8	1-8
spiny	C	8	1-8
Poinsettia, wild	C	-	-
Puncturevine	C	-	-
Purslane, common	C	-	-



**Manufactured for: Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**BROADLEAF WEEDS CONTROLLED (continued)**

Broadleaf Weeds	SOIL APPLIED	POST-EMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Pusley, Florida	C	-	-
Ragweed, Common Giant	R R	R R	1-3 1-3
Sage, barnyard	R	1-3	-
Sida, prickly	C*	-	-
Smartweed, ladysthumb Pennsylvania	C C	4 4	1-3 1-3
Spurge prostrate spotted	C C	4 4	1-3 1-3
Sunflower	C*	4	1-3
Thistle, Canada	-	R	1-3
Velvetleaf	C*	4	1-3

\*When **Tigris Imazethapyr 2SL** is soil applied, these weeds are more consistently controlled by pre-plant incorporated treatments.

**GRASS WEEDS CONTROLLED\***

Broadleaf Weeds	SOIL APPLIED	POST-EMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Barnyardgrass	R	3	1-3
Crabgrass, large smooth	R R	3 3	1-3 1-3
Cupgrass, woolly***	-	3	1-3
Foxtail, giant green yellow	C C C	6 3 3	1-6 1-3 1-3
Goosegrass	R	-	-
Johnsongrass, seedling rhizome	R C	6 R	1-8 6-12
Panicum, fall Texas	R R	- -	- -
Red rice	-	3	1-3
Shattercane	R	6	1-8
Signalgrass, broadleaf	R	4	1-8
Sorghum, alnum	R	6	1-3

\*Pre-plant incorporated treatments of **Tigris Imazethapyr 2SL** are more consistent for grass control.

\*\***Tigris Imazethapyr 2SL** is active against many broadleaf and grass species. However, when heavy grass or common lambsquarters pressure is anticipated, use **Tigris Imazethapyr 2SL** in combination with a registered soil-applied grass herbicide (such as pendimethalin) for optimum control (See **HERBICIDE COMBINATIONS** section).

\*\*\***Tigris Imazethapyr 2SL** only controls emerged woolly cupgrass.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557



## SEDGES

Weeds Controlled	SOIL APPLIED	POST-EMERGENCE	
		Maximum Leaf Stage	Size (Inches)
Nutsedge, purple yellow	R	R	1-3
	R	R	1-3

## HERBICIDE COMBINATIONS

### GRASS WEEDS

Use a soil applied grass herbicide (such as pendimethalin) to control grass weeds not on the **Tigris Imazethapyr 2SL** label and to enhance the control of certain broadleaf weeds such as common lambsquarters and pigweeds. Refer to the pendimethalin (or other grass herbicide) label for specific use instructions, rates and precautions.

When applied as directed, **Tigris Imazethapyr 2SL** pre-plant incorporated or pre-emergence combination treatments with pendimethalin, trifluralin, dimethenamide-P, alachlor, or S-metholachlor will control the weeds listed in following table, in addition to those controlled by **Tigris Imazethapyr 2SL** alone.

GRASSES	Pendimethalin <sup>a</sup>	Trifluralin <sup>b</sup>	Alachlor	S-metholachlor	Dimethenamide-P
Barnyardgrass	X	X	X	X	X
Crabgrass, smooth	X	X	X	X	X
Crabgrass, large	X	X	X	X	X
Crowfootgrass	X	X			
Goosegrass	X	X	X	X	X
Millet, wild proso	X	X			
Panicum, fall	X	X	X	X	X
Panicum, Texas	X	X			
Sandbur, field	X	X			
Shattercane	X <sup>b</sup>	X			
Signalgrass, broadleaf	X <sup>b</sup>	X	X	X	X
Witchgrass	X	X	X	X	X

<sup>a</sup> Pre-plant incorporated tank-mixture applications of **Tigris Imazethapyr 2SL** plus pendimethalin will suppress the growth of itchgrass and rhizome johnsongrass.

<sup>b</sup> Pre-plant incorporated treatments only

A selective post-emergence grass herbicide such as sethoxydim may be mixed with **Tigris Imazethapyr 2SL** to control volunteer corn or grasses not controlled by **Tigris Imazethapyr 2SL**. For best results use crop oil concentrate AND liquid fertilizer with grass herbicide tank-mixtures.



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

### Tigris Imazethapyr 2SL+ Sethoxydim For Enhanced Grass Control

Apply **Tigris Imazethapyr 2SL** at the rate of 4 fl. oz./Acre (0.063 lb. ai/Acre). The addition of sethoxydim to **Tigris Imazethapyr 2SL** at the specified rates will control the grasses listed below. (Refer to the sethoxydim label for additional weeds controlled.)

Sethoxydim (oz. per acre)	Annual Grasses Controlled	Size (inches)
Refer to product label	Wild Proso Millet	4 - 10"
	Shattercane	3 - 12"
	Foxtail, Giant	3 - 8"
	Junglerice	3 - 8"
	Panicum, Fall Texas	3 - 8" 3 - 8"
	Signalgrass, Broadleaf	3 - 8"
	Volunteer Corn	4 - 10"
	Barnyardgrass	3 - 8"
	Crabgrass, Large Smooth	3 - 6" 3 - 6"
	Cupgrass, Woolly	3 - 8"
	Foxtail, Green Yellow	3 - 8" 3 - 8"
	Goosegrass	3 - 6"
	Johnsongrass, Seedling	3 - 8"
	Sprangletop, Red	3 - 8"
	Witchgrass	3 - 8"

\*If a mixture of grasses are present, use the highest rate indicated for the grasses present.

The addition of sethoxydim to **Tigris Imazethapyr 2SL** enhances the grass control, especially when heavy infestations of grass exist. It also provides control of grasses not controlled **Tigris Imazethapyr 2SL**. In some cases the activity of sethoxydim may be reduced when mixed with **Tigris Imazethapyr 2SL**. The reduction in activity may be overcome by delaying the application of sethoxydim herbicide 7 days following the application of **Tigris Imazethapyr 2SL**. If sethoxydim is applied first, wait 3 days before applying **Tigris Imazethapyr 2SL**.

For optimum control, apply the tank-mixture to actively growing weeds at the sizes indicated in the table above (for sequential applications refer to application rates and weeds sizes indicated in the **Tigris Imazethapyr 2SL** and sethoxydim labels). Refer to the sethoxydim label for additional information regarding application rates, restrictions, precautions, weeds controlled, adjuvants specified and other information.

#### BROADLEAF WEEDS

Broadleaf herbicides that can be tank-mixed with **Tigris Imazethapyr 2SL** include acifluorfen, sodium bentazon, lactofen, cloransulam-methyl, paraquat dichloride, sodium bentazon + sodium acifluorfen, or fomesafen. Glyphosate may be tank-mixed with **Tigris Imazethapyr 2SL** to aid in control of certain weeds only in Roundup Ready® Soybeans. See the glyphosate label for rates and weeds controlled and other restrictions. DO NOT apply **Tigris Imazethapyr 2SL** with certain herbicides (see RESTRICTIONS section).

#### **Tigris Imazethapyr 2SL+ Acifluorfen For Enhanced Control of Common Ragweed and Pigweeds (including tall and common waterhemp)**

The addition of acifluorfen to **Tigris Imazethapyr 2SL** at the specified rates will enhance the control of several broadleaf weeds, including common and giant ragweed, pigweed species and waterhemp. (Refer to the acifluorfen label for additional weeds controlled.)

When tank-mixing acifluorfen with **Tigris Imazethapyr 2SL**, apply **Tigris Imazethapyr 2SL** at the rate of 4 fl. oz./Acre (0.063 lb. ai/Acre). Refer to the acifluorfen label for applicable rates and directions for use.

#### **Tigris Imazethapyr 2SL+ Cloransulam-methyl for Enhanced Control of Ragweed Species**

Cloransulam-methyl may be tank-mixed with **Tigris Imazethapyr 2SL** to aid in the control of common and giant ragweed. See the cloransulam-methyl label for specified rates and precautions.

#### **Tigris Imazethapyr 2SL+ Sulfentrazone Containing Compounds**

**Tigris Imazethapyr 2SL** provides control of many grasses and broadleaf weeds when applied to the soil or applied post-emergence to weeds. It also provides season-long control of many weeds. Sulfentrazone containing products may be tank-mixed with **Tigris Imazethapyr 2SL** in soil applications for enhanced weed control in soybeans.

**Tigris Imazethapyr 2SL** may be applied post-emergence to soybeans previously treated with sulfentrazone containing products.

NOTE: Sulfentrazone-containing products are only labeled for soil applications to soybean..



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

### **Tigris Imazethapyr 2SL+ Thifensulfuron for Enhanced Control of Common Lambsquarters**

For optimal weed control management, apply a soil applied grass herbicide such as pendimethalin or trifluralin followed by **Tigris Imazethapyr 2SL** post-emergence. If common lambsquarters are not adequately controlled by the soil applied treatment, thifensulfuron may be tank-mixed with **Tigris Imazethapyr 2SL** for additional activity.

The addition of thifensulfuron to **Tigris Imazethapyr 2SL** may cause severe injury and/or stunting to soybeans, especially when applied under hot, humid conditions. The USER ASSUMES ALL RISKS AND CONSEQUENCES associated with applications of this tank-mixture to soybeans.

When tank-mixing thifensulfuron with **Tigris Imazethapyr 2SL**, use the following rates:

**Tigris Imazethapyr 2SL**– 4 fl. oz./Acre (0.063 lb. ai/Acre)

AND

Specified label rate of thifensulfuron

Add to the spray mixture:

Non-ionic surfactant – 1 quart per 100 gals. (0.25% v/v)

AND

Liquid nitrogen based fertilizer (such as 28%N, 32%N, or 10-34-0) at the rate of 1.25 to 2.5 gals. per 100 gals. of spray solution. Instead of a liquid fertilizer, spray grade ammonium sulfate may be used at the rate of 12 – 15 lbs. per 100 gals. of spray solution.

Apply to 1 – 3 trifoliolate stage soybeans only.

### **Other Tank-Mixture Combinations**

#### **Tigris Imazethapyr 2SL+ Imazaquin for Volunteer Corn and Common Sunflower**

The application of **Tigris Imazethapyr 2SL** plus imazaquin may be applied to states or portions of states described as Region 2 or Region 3 on the imazaquin label, and the following counties in South Dakota: Yankton, Bon Homme, Hutchinson, McCook, Hanson, Davison, Miner, Lake, and Kingsbury. Refer to the respective labels for the specified use area. DO NOT use this tank-mixture in North Dakota or in Minnesota north of state highway #210.

Apply the products at the following rate:

**Tigris Imazethapyr 2SL**– 4 fl. oz./Acre (0.063 lb. ai/A)

AND

Specified label rate of imazaquin

The tank-mixture of **Tigris Imazethapyr 2SL** plus imazaquin will suppress volunteer corn. Apply to volunteer corn up to 10 inches in height.

The tank-mixture of **Tigris Imazethapyr 2SL** and imazaquin will enhance the control of common sunflowers. Apply to sunflowers up to 3 inches in size.

Refer to the imazaquin label for additional weeds controlled.

A post-emergence application of **Tigris Imazethapyr 2SL** plus imazaquin will NOT suppress volunteer CLEARFIELD® corn (field corn hybrids which possess resistance to imidazolinone herbicides i.e., **Tigris Imazethapyr 2SL** and imazaquin).

### **APPLICATIONS TO SOYBEANS IN NORTH DAKOTA AND MINNESOTA (north of highway #210)**

**Application Rate:** Apply **Tigris Imazethapyr 2SL** at 3 fl. oz./Acre (0.047 lb. ai/Acre) post-emergence only.

Weeds Controlled	POST-EMERGENCE	
	Maximum Leaf Stage	Size (Inches)
Cocklebur, common*	4	1 – 4
Kochia (non-ALS resistant)	4	1 – 3
Mustard sp.	4	1 – 3
Nightshade, black	4	1 – 3
Eastern black	4	1 – 3
hairy	4	1 – 3
Pigweed, redroot	4	1 – 4
Wild oats**	3	1 – 4

\*For control of common cocklebur, add acifluorfen at the label specified rate to the spray solution.

\*\***Tigris Imazethapyr 2SL** will reduce competition from wild oats.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying **Tigris Imazethapyr 2SL** at the specified rate: (Planting earlier than the specified interval may result in crop injury.)

Crop	Months after Tigris Imazethapyr 2SL application
CLEARFIELD® corn hybrids (resistant to <b>Tigris Imazethapyr 2SL</b> ) Lima beans Southern peas Soybeans Peanuts	Anytime
Alfalfa Clover Rye (Except in North Dakota and Minnesota north of highway #210) Wheat Edible beans and peas (other than lima beans and Southern peas)	4
Field corn Field corn grown for seed	8 1/2
Barley Tobacco	9 1/2
Cotton* Lettuce Oats Popcorn Rye in North Dakota and Minnesota north of highway #210 Safflower Sorghum Sunflower Sweet corn	18
Potatoes Flax	26
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE**	40

\*Refer to the following table for a Cotton Rotation Interval following **Tigris Imazethapyr 2SL** application to alfalfa or clover grown for seed production. These guidelines do not apply to **Tigris Imazethapyr 2SL** applications made to alfalfa grown for hay or forage (Use the 18-month Rotational Interval above).

\*\*Following forty months after a **Tigris Imazethapyr 2SL** application, and before planting any crop not listed elsewhere in the ROTATIONAL CROP RESTRICTIONS, a successful field bioassay must be completed. The field bioassay consists of a test strip of the intended rotational crop planted across the previously treated field and grown to maturity. The test strip must include low areas and knolls, and include variations in soil such as type and pH. If no crop injury is evident in the test strip, the intended rotational crop may be planted the following year.

Sugar beet production can be reduced when grown in soil conditions with a pH less than 6.5. If the field is limed to adjust pH prior to planting rotational crops not listed in the ROTATIONAL CROP RESTRICTIONS, apply the lime at least 12 months prior to planting the rotational crop.

Use of **Tigris Imazethapyr 2SL** herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

### Cotton Rotation Following Application of Tigris Imazethapyr 2SL to Alfalfa or Clover Grown For Seed

Irrigation/Precipitation Requirements	Rotation Interval	
	Less than 3 acre feet or 36" of water	40 Months
	Greater than or equal to 3 acre feet or 36" of water	18 Months

### EXCEPTIONS TO ROTATIONAL CROP RESTRICTIONS

**Barley:** (States of Delaware, Indiana, Kentucky, Maryland, New Jersey, Ohio, Pennsylvania, and Virginia only.) Barley may be planted 4 months following a **Tigris Imazethapyr 2SL** application in these states.

**Barley:** (North Dakota only.) Barley may be planted 18 months following an **Tigris Imazethapyr 2SL** application.

**CLEARFIELD® canola:** CLEARFIELD® varieties of canola, such as Pioneer® 45A71 and Pioneer® 46A76, may be planted as a rotational crop the next season after an application of **Tigris Imazethapyr 2SL** herbicide at label rates on registered crops.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

**Corn inbred lines:** Corn inbred seed lines may be planted the year following an application of **Tigris Imazethapyr 2SL**. Several seed companies have tested a wide range of inbreds for sensitivity to **Tigris Imazethapyr 2SL** soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, Tigris, LLC has not been given access to the inbred data. Growers are directed to contact the seed company for information and instructions regarding the planting of corn grown for seed in fields treated with **Tigris Imazethapyr 2SL** the previous year. Since growing conditions, environmental conditions and grower practices are beyond the control of Tigris, LLC, all risks and consequences associated with planting seed corn inbreds into fields treated previously with **Tigris Imazethapyr 2SL** shall, to the extent allowable by applicable law, be assumed by the user.

**Sweet corn and popcorn varieties:** (States of Illinois, Indiana, Iowa, Minnesota, Ohio, Tennessee, and Wisconsin only.) Sweet corn and popcorn varieties may be planted the year following an application of **Tigris Imazethapyr 2SL**. Some sweet corn and popcorn varieties may be injured when planted at less than 18 months following an application of **Tigris Imazethapyr 2SL** herbicide. Before planting sweet corn for processing, contact the processor company for information and specifications regarding the sensitivity of sweet corn varieties planned for fields treated with **Tigris Imazethapyr 2SL** the previous year. **DO NOT** plant fresh market sweet corn varieties prior to 18 months after **Tigris Imazethapyr 2SL** use. Before planting popcorn, contact the popcorn company for information and instructions regarding the sensitivity of popcorn varieties planned for fields treated with **Tigris Imazethapyr 2SL** the previous year.

Since growing conditions, environmental conditions and grower practices are beyond the control of TIGRIS, LLC, TO THE EXTENT ALLOWABLE BY APPLICABLE LAW, ALL RISKS AND CONSEQUENCES ASSOCIATED WITH PLANTING SWEET CORN OR POPCORN VARIETIES INTO FIELDS TREATED PREVIOUSLY WITH **Tigris Imazethapyr 2SL** SHALL BE ASSUMED BY THE USER.

Stunting and maturity delay or other adverse effects may result when sweet corn or popcorn are planted following **Tigris Imazethapyr 2SL** use.

**Certain vegetable crops:** (States of Alabama, Delaware, Florida, Georgia, Indiana, Kentucky, Maryland, New Jersey, North Carolina, Pennsylvania, South Carolina, and Virginia only.) The following crops may be planted 18 months following the last application of **Tigris Imazethapyr 2SL**: bahiagrass, cabbage, cantaloupe, cucumber, Irish potato, onion, sweet potato transplants, sweet pepper transplants, tomato transplants, and watermelon.

**Cotton:** (States of North Carolina, South Carolina, and Virginia only.) Cotton may be planted nine and one-half months after an application of **Tigris Imazethapyr 2SL** if all of the following criteria are met:

- **Tigris Imazethapyr 2SL** applied to peanuts only.
- Soil texture is sandy loam or loamy sand only.
- Greater than 16 inches of rainfall and/or irrigation is received following application of **Tigris Imazethapyr 2SL** through October of the application year.

**Field Corn and Field Corn Grown for Seed:** (Arizona, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming) Nine and one-half months after **Tigris Imazethapyr 2SL** application.

**Snap Beans:** When applied at no more than 1.5 fl. oz./Acre (0.023 lb. ai/Acre) to snap beans in the use areas defined on this label, snap beans may be replanted at any time after application of **Tigris Imazethapyr 2SL**.

**Wheat:** Wheat may be planted 3 months following an **Tigris Imazethapyr 2SL** application in areas east of Interstate highway I-35.

When **Tigris Imazethapyr 2SL** is applied at no more than 3 oz./Acre (0.047 lb. ai/A) to edible legumes in the use areas described the following rotational restrictions apply:

- Chickpeas, lentils and peas may be planted anytime following an **Tigris Imazethapyr 2SL** application.
- Snap beans may be planted 3 months and barley 4 months following an application of **Tigris Imazethapyr 2SL**.

#### ADDITIONAL USE INFORMATION

##### EDIBLE LEGUMES VEGETABLES

###### Restrictions:

- There must be an interval of at least 30 days between application and harvest of snap beans, lima beans, chickpeas (Arizona and California), English peas, and Southern peas.
- There must be an interval of at least 60 days between application and harvest of dry edible peas, lentils, chickpeas, red kidney beans, and other dry bean or pea types listed on this label.

When **Tigris Imazethapyr 2SL** is applied at no more than 3 fl. oz./Acre (0.047 lb. ai/A) to edible legumes in the use areas described, the following rotational restrictions apply:

- Chickpeas, lentils and peas may be planted anytime following an **Tigris Imazethapyr 2SL** application.
- Snap beans may be planted 3 months and barley 4 months following an application of **Tigris Imazethapyr 2SL**.

##### NON-GRASS ANIMAL FEED

###### (ALFALFA AND CLOVER)

**Restriction:** DO NOT feed, graze or harvest alfalfa or clover for 30 days following an application of **Tigris Imazethapyr 2SL** to alfalfa or clover.

##### SOYBEANS

###### Restrictions:

- There must be an interval of at least 85 days between an application of **Tigris Imazethapyr 2SL** and soybean harvest.
- Make **Tigris Imazethapyr 2SL** applications before soybean bloom.
- DO NOT graze or feed treated soybean forage, hay or straw to livestock.
- DO NOT tank-mix **Tigris Imazethapyr 2SL** with clomazone containing herbicides. **Tigris Imazethapyr 2SL** may be applied post-emergence following a soil application of clomazone.

If soybeans are furrow irrigated, till the soil prior to planting winter wheat or barley. Break the beds up and the soil mixed with tillage equipment set to cut 4 to 6 inches deep.



Manufactured for: **Tigris, LLC**

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557

## PEANUTS

### Restrictions:

- Do not graze or feed treated peanut forage, vines, hay or straw to livestock.
- There must be an interval of at least 85 days between an application of **Tigris Imazethapyr 2SL** and peanut harvest.
- DO NOT apply pendimethalin + imazethapyr herbicides to peanuts the same year as **Tigris Imazethapyr 2SL**.

Chlorimuron may be applied post-emergence to peanuts following a **Tigris Imazethapyr 2SL** application. Refer to the chlorimuron label for specific use directions.

### ALL CROPS

Full rate application of products containing chlorimuron ethyl, chloransulam-methyl, flumetsulam, imazaquin, or products containing imazethapyr the same year as **Tigris Imazethapyr 2SL** may increase the risk of injury to sensitive follow crops. Consult labels for specific uses of these products in combinations.

Only rotational crops harvested at maturity may be used for feed or food.

In the event of a crop loss due to weather, soybeans, peanuts, or CLEARFIELD® corn can be replanted. DO NOT work the soil deeper than 2 inches.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep from freezing: DO NOT store below 32°F.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA region office for guidance.

### CONTAINER HANDLING:

**NONREFILLABLE CONTAINER (EQUAL TO OR LESS THAN 5 GALLONS):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

**NONREFILLABLE CONTAINER (GREATER THAN 5 GALLONS):** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available.

**REFILLABLE CONTAINER:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Tigris, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Tigris, LLC and Seller harmless for any claims relating to such factors.

Tigris, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or Tigris, LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, Tigris, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither Tigris, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF TIGRIS, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF TIGRIS, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Tigris, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Tigris, LLC.

Clearfield is a registered trademark of BASF.

Pioneer is a registered trademark of Agco Corporation.

[EPA 20190813]

2020.11.06V1



Manufactured for: Tigris, LLC

P.O. Box 250

10025 Hwy. 264 Alternate

Middlesex, NC 27557