

For years, growers and ranchers have counted on WeedMaster® for consistent weed control in sugarcane, pastures, small grains and fallowland. WeedMaster is a selective, post-emerge herbicide that provides tough, long-lasting control of a broad spectrum of annual, biennial and perennial broadleaf weeds. The growth regulating effects of 2,4-D and dicamba interfere with the normal development of weeds, ultimately causing plant death.

COMPETITIVE ADVANTAGES

CONSISTENT CONTROL IN A CONCENTRATED FORMULA

- Achieve outstanding control of brush and broadleaf weeds in a concentrated formula
- Controls tough weeds such as horsenettle, thistles and blackberry
- Safe for desirable grasses
- Pre-emergence control with residual activity

AT-A-GLANCE INFO

EPA REG. NO.	71368-34
ACTIVE INGR.	DMA dicamba salt (12.4%), 2,4-D amine salt (35.7%)
FORMULATION	liquid
CHEM. FAMILY	benzoic acid, phenoxycarboxylic acid
HRAC NO.	4
MODE OF ACT	auxin growth regulator
REI	48 hours
SIGNAL WORD	danger
RESTRICTED	no
PACKAGE SIZE	1 gal container (5 per case), 2.5 gal container (2 per case), 30 gal drum,

265 gal tote, bulk









DEPEND ON WEEDMASTER®.

KEY USES

Conservation Reserve Program (CRP) land

Fallowland

General farmstead

Sorghum

Grass (hay or silage)

Pasture Rangeland

Sugarcane

Wheat

KEY WEEDS CONTROLLED

(See product label for complete list)

ANNUALS BIENNIALS & Buckwheat, wild **PERENNIALS** Burdock Bindweed, field Buttercup **Bittercress** Cockle, cow Chicory Cocklebur, Dandelion, common common Dogfennel Dock, curly

Evening primrose Goldenrod, Missouri Fleabane, annual Groundsel, Texas Honeysuckle, hairy Knotweed Horsenettle, Kochia Carolina

Lambsquarters, Ivy, poison
common Marshelder
Marestail Mesquite
(horseweed) Persimmon, eastern

Morningglory; Prickly lettuce ivyleaf, tall Smartweed Mustards; annual, tansy Thistle, bull Pigweed; Prickly lettuce Smartweed Smartweed Thistle, bull Vetch, hairy

prostrate, redroot, smooth, tumble Purslane, common Ragweed; common, lanceleaf, western

Sowthistle

Shepherd's-purse Smartweed,

Pennsylvania

Sneezeweed, bitter

Thistle, Russian Velvetleaf

USE SUMMARY

RATES: 0.5 pts per acre to 4 to 5 pts/A for control of annual, biennial and perennial weeds. See label for per acre rate according to weed growth stage.

Apply to actively growing weeds as aerial, broadcast, band, or spot spray applications.

Apply with water or sprayable fluid fertilizer as a carrier. Sprayable fluid fertilizer may be used as the carrier in preplant or pre-emergence uses. Post-emergence uses with sprayable fluid fertilizer may be made only on pasture, hayland, or wheat crops.

Surfactants and/or low use rate liquid fertilizers (28-0-0, 32-0-0) or crop oil concentrate may be used with this product to improve burndown of emerged weeds. (Do not use crop oil concentrate for feed crops – refer to crop oil concentrate label for complete directions.)

AERIAL APPLICATION: Use 3 to 10 gals of water/A.

GROUND APPLICATION (BROADCAST): Use 5 to 40 gals per broadcast acre.

See label for complete directions for use.



