



Nufarm

# Weedone® LV6 EC

SELECTIVE HERBICIDE

## WEED CONTROL WITHOUT INJURY TO ESTABLISHED GRASSES.

Weedone® LV6 EC offers easy handling and storing and is ideal for aerial and custom applications. Mixes with liquid fertilizer and can also be tank mixed with water and oil for difficult to control perennial broadleaf weeds and brush.

### COMPETITIVE ADVANTAGES

#### FLEXIBLE MIXING AND APPLICATION OPTIONS

- Low volatile formulation mixes with liquid fertilizers
- Mixes with water and oil to control tough perennial broadleaf weeds and brush
- Highest acid concentration, allows for efficient shipping and storage
- Ideal for aerial application
- Compatible with Diablo® and other products for complete kochia control

### AT-A-GLANCE INFO

EPA REG. NO.	71368-11
ACTIVE INGR.	2,4-D ester (87.3%)
FORMULATION	liquid
CHEM. FAMILY	phenoxy-carboxylic acid
GROUP NO.	4
MODE OF ACT.	synthetic auxin
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	2.5 gal container (2 per case), 30 gal

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

 BurnMaster

 Panther SC

 WeedMaster®

 Nufarm  
Grow a better tomorrow

# Weedone® LV6 EC

## WEED CONTROL WITHOUT INJURY TO ESTABLISHED GRASSES.

### KEY USES

Fencerows, roadsides, vacant lots, airfields, railroads, highway and utility rights-of-way, pastures, rangelands, forest management, and other non-crop areas

### KEY WEEDS CONTROLLED

(See product label for complete list)

- Alfalfa
- Bull thistle
- Chicory
- Cocklebur
- Dandelion
- Knotweed
- Kochia
- Lambsquarters
- Marshelder
- Mustard
- Pigweed
- Ragweed
- Russian thistle
- Sowthistle
- Velvetleaf
- Wild garlic
- Wild lettuce
- Wild onion

### USE SUMMARY

#### RANGELAND AND PASTURES

For ground application, a minimum of 10–15 gallons of water per acre is recommended for proper spray coverage.

For aerial application, use a minimum of 2 gallons of water per acre.

Do not graze lactating dairy animals on treated areas within 7 days after treatment.

Do not cut grass for hay within 30 days after application.

Withdraw meat animals being finished for slaughter from treated forage at least 3 days before slaughter.

#### NON-CROP AREAS

For aerial application to solid stands of susceptible brush, use 1-1/3 to 2-2/3 quarts in 3–12 gallons volume per acre. The use of a high quality adjuvant or oil is permissible.

See product label for complete directions for use.

#### APPLICATION RATES

##### Weeds and brush in pastures and rangeland

Annual Broadleaf Weeds.....	21 fl oz/A
Perennial and Biennial Broadleaf Weeds.....	32 to 43 fl oz/A

##### Fencerows, roadsides, vacant lots, airfields, railroad, highway and utility rights-of-way and other non-crop areas

Annual Broadleaf Weeds.....	21 to 43 fl oz/A
Perennial and Biennial Broadleaf Weeds.....	32 to 64 fl oz/A
Cattails and Tules.....	16 to 27 fl oz/A
Woody Plants.....	11 to 21 fl oz/A

See product label for complete application rates.

