

# Chick Brooder - Weekly Capacity

*Newborns require heat for 4 - 6 weeks*

Highlighted capacities indicate the max number of weeks heat is required from brooder

## 9.5" Chick Brooders (Short) (Model: CB25)

Quail	Week	2 Section	1 Layer	2 Layer	3 Layer	4 Layer	5 Layer
	1	30-35	45-55	90-110	135-165	180-220	225-275
	2	25-30	35-45	70-90	105-135	140-180	175-225
	3	20-25	30-40	60-80	90-120	120-160	150-200
	4	16-20	24-30	48-60	72-90	96-120	120-150

Pheasant Partridge	Week	2 Section	1 Layer	2 Layer	3 Layer	4 Layer	5 Layer
	1	25-30	35-45	70-90	105-135	140-180	175-225
	2	20-25	30-35	60-70	90-105	120-140	150-175
	3	18-20	25-30	50-60	75-90	100-120	125-150
	4	14-18	20-25	40-50	60-75	80-100	100-125
5	12-14	18-21	36-42	54-63	72-84	90-105	

## 15" Chick Brooders (Tall) (Model: CB40)

Chicken Duck	Week	2 Section	1 Layer	2 Layer	3 Layer	4 Layer
	1	20-22	30-35	60-70	90-105	120-140
	2	18-20	25-30	50-60	75-90	100-120
	3	15-18	20-25	40-50	60-75	80-100
	4	12-14	18-21	36-42	54-63	72-84
	5	10-12	15-18	30-36	45-54	60-72
6	8-10	12-15	24-30	36-45	48-60	

Turkey Goose	Week	2 Section	1 Layer	2 Layer	3 Layer	4 Layer
	1	10-12	15-18	30-36	45-54	60-72
	2	8-10	12-15	24-30	36-45	48-60
	3	7-9	10-13	20-26	30-39	40-52
	4	6-8	9-12	18-24	27-36	36-48
	5	5-7	8-10	16-20	24-30	32-40
6	4-6	6-9	12-18	18-27	24-36	

*After 4 - 6 weeks, move the birds to a grow-out pen, breeding cage, or free-range.*