

# TexaClear® Congestion and Sinus Pain Relief 8oz

## Drug Facts

Active ingredients (in each 30 mL)	Purpose
Choline Salicylate (NSAID)* 870 mg.....	Pain reliever/fever reducer
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

\*nonsteroidal anti-inflammatory drug

## Uses

- temporarily relieves
  - nasal congestion
  - headache
  - minor aches and pains
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid bronchial passageways of bothersome mucus and make coughs more productive

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.

When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Salicylates (NSAIDs) may cause a severe allergic reaction which may include:

- hives
- rash
- shock
- skin reddening
- facial swelling
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Do not use

- if you have ever had an allergic reaction to salicylates (including aspirin) or any other pain reliever/fever reducer.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

## Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn or if you have ulcers or bleeding problems
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, diabetes or thyroid disease
- you have difficulty in urination due to enlargement of the prostate gland
- you have a cough with excessive phlegm (mucus) ▶

## Drug Facts (continued)

- you have a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- you are taking a diuretic
- you have asthma

## Ask a doctor or pharmacist before use if you are

- taking a prescription drug for anticoagulation (thinning the blood) diabetes, gout, or arthritis.
- under a doctor's care for any serious condition
- taking a diuretic.

## When using this product do not use more than directed

## Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use salicylates (NSAID) during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- do not exceed recommended dosage.
- do not take more than 6 doses in any 24-hour period.
- use enclosed dose cup and keep with product
- mL= milliliter
- adults and children 12 years of age and over: 30 mL every 4 hours
- children under 12 years of age: do not use

## Other information

- each 30mL contains: **Sodium 17 mg**
- read all product information before using
- store at 20° to 30°C (68° to 86°F)

## Inactive ingredients

citric acid anhydrous, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol

## Questions or comments?

Call 1-888-535-0305 9 a.m. - 5 p.m. CST

**GM Pharmaceuticals, Inc.**

Distributed by: GM Pharmaceuticals, Inc. Arlington, TX 76015