Water breaks, active straining, foal emerges

Foal to nurse within 2 hours

Passing of placenta

Mare to pass placenta within 3 hours

Up to 33% of Mares Experience Complications or Loss Before, During, and/or After Birth

BEFORE

1/3

of mares are at risk of losing their foal before birth (most within first 35 days)

DURING

10%

of mares experience complications during foaling (dystocia most common)

AFTER

15%

of mares experience serious complications after foaling (for up to 3 months)

Complications and Compromised Mare Health Increases Odds of Foal Death

Foals from mares with compromised health have greater odds of death

- 19x Abnormal placenta
- **18x** History of illness during pregnancy



Foals with complications are more likely to die

- 21x Foals with seizures (in the first 24 hours)
- **16x** Foals with complications (during hospitalizaton)



Be notified to early signs of foaling with a smart halter™

so you can be by their side when they need you most, and closely monitor your mare for complications before, during, and after birth.



Place your smart halter™ on your pregnant mare

30 days prior to her foal date and monitor her for at least 3 months thereafter.



A single smart halter™ can be transferred

between multiple horses making it a must have for every owner, breeder, trainer, and veterinarian this foaling season.



The NIGHTWATCH® smart halter™ is an equine distress & wellness monitor that offers automated alerts and objective insight for your performance, pregnant, and high-risk horses. From single horse owners and veterinary clinics to large multi-national breeding and training programs, NIGHTWATCH® is a tool that can add value to equine operations of all sizes.