

FOR ANIMAL USE ONLY

TRIWORM-D

Reg. No. G3645 Act 36/1947

INDICATIONS:

Worm remedy against ascarids, hookworms, whipworms and tapeworms in dogs.

COMPOSITION:

Each tablet contains:

Praziquantel..... 50 mg
Pyrantel pamoate 144 mg
Febantel 150 mg

STORAGE INSTRUCTIONS:

Store below 25 °C.

WARNINGS:

- DO NOT USE SIMULTANEOUSLY WITH PIPERAZINE COMPOUNDS.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE:

USE ONLY AS DIRECTED.

- Can be administered directly, or mixed with a small amount of food, or crushed and mixed with feed.
- Feeding or fasting before treatment is not necessary.

Dosage:

Dose according to body mass i.e. 1 tablet/10 kg

Body mass (kg)	Dosage
0,5 – 2	¼ tablet
2 – 5	½ tablet
6 – 10	1 tablet
11 – 15	1 ½ tablets
16 – 20	2 tablets
21 – 25	2 ½ tablets
26 – 30	3 tablets
31 – 35	3 ½ tablets
36 – 40	4 tablets
40 – 50	5 tablets

- Puppies should be dewormed from the age of 2 – 12 weeks at 2 week intervals, and at regular intervals of 3 – 6 months thereafter to control worm infestation.
- Triworm-D can be given safely to all dogs including young pups, old dogs, working dogs and pregnant bitches 10 days before whelping.

EFFICACY:

Controls:

Ascarids	<i>Toxocara canis</i>
Hookworms	<i>Ancylostoma caninum</i>
Tapeworms	<i>Dipylidium caninum</i> ; <i>Taenia</i> spp. <i>Echinococcus</i> spp.
Whipworm	<i>Trichuris vulpis</i>

Registration holder:

Cipla Vet (Pty) Limited
Co. Reg. No. 2001/017471/07
P O Box 1096, Durbanville, 7551
Tel: 0861 115 037

xxxx x

SLEGS VIR DIEREGERBUIK

TRIWORM-D

Reg. Nr. G3645 Wet 36/1947

AANWYSINGS:

Wurmmiddel teen spoelwurms, haakwurms, sambokwurms en lintwurms in honde.

SAMESTELLING:

Elke tablet bevat:

Prasikwantel 50 mg
Pirantel pamoaat 144 mg
Febantel 150 mg

BERGINGSANWYSINGS:

Berg onder 25 °C.

WAARSKUWINGS:

- MOENIE SAAM MET PIPERASIENBEVATTENDE MIDDELS GEBRUIK NIE.
- Hou buite bereik van oningeligte persone, kinders en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEBRUIKSAANWYSINGS:

GEBRUIK SLEGS SOOS AANGEDUI

- Kan direk toegedien word, óf met 'n bietjie vleis gemeng, óf verkruimel en met kos gemeng.
- Dit is nie nodig om diere te voer of uit te honger voor toediening nie.

Dosis:

Doseer volgens liggaamsmassa d.w.s. 1 tablet/10 kg

Liggaamsmassa (kg)	Dosis
0,5 – 2	¼ tablet
2 – 5	½ tablet
6 – 10	1 tablet
11 – 15	1 ½ tablette
16 – 20	2 tablette
21 – 25	2 ½ tablette
26 – 30	3 tablette
31 – 35	3 ½ tablette
36 – 40	4 tablette
40 – 50	5 tablette

- Ontwurm jong hondjies 2-weekliks vanaf 2 - 12 weke, en daarna met tussenposes van 3 - 6 maande om wurmbesmetting te beheer.
- Kan met veiligheid toegedien word aan jong hondjies, ou honde, werkende honde, en dragtige tewe tot 10 dae voor geboorte.

DOELTREFFENDHEID:

Beheer:

Spoelwurms	<i>Toxocara canis</i>
Haakwurms	<i>Ancylostoma caninum</i>
Lintwurms	<i>Dipylidium caninum</i> ; <i>Taenia</i> spp. <i>Echinococcus</i> spp.
Sambokwurm	<i>Trichuris vulpis</i>

Registrasiehouer:

Cipla Vet (Edms) Bpk
Mpy. Reg. Nr. 2001/017471/07
Parc du Cap, Building 9, Mispel Street,
Belville, 7530, RSA
Tel: 0861 115 037

xxxx x