

Minimizing the Pain of Chemo and Radiation Side Effects

For Prevention of Radiation Dermatitis

When to Use RadiaPlex[®] Gel:

- Prior to radiation and radiation dermatitis
- Dry & moist desquamation skin
- 1st & 2nd Degree Burns
- Cuts & Abrasions



(Prescription)

How to Use:

- Cleanse with **MPM Wound and Skin Cleanser** and allow drying
- Apply **RadiaPlex Gel** 1/4" thick – 3 times daily during treatment plus 2 weeks after treatment
- Cleanse the irradiated area with a suitable wound cleanser such as MPM Wound & Skin Cleanser.
- Apply a generous amount of RadiaPlex from the first day of treatment three times per day, seven days a week to the treated areas; gently rub/message into the areas until RadiaPlex is completely absorbed.
- RadiaPlex should be applied immediately after treatment 3 times a day for maximum results.
- RadiaPlex may be applied as indicated by the treating physician. Do not apply 2 hours before treatment or directed by the treating physician.
- Continue to apply RadiaPlex as described above until the skin has fully recovered.
- Do not interrupt applications during the course of the radiation therapy, even for one day.

Benefits to Patient:

- Adheres to skin tissue
- Provides moisture to skin cells
- Helps prevent radiation damage
- Non greasy or sticky

For Treatment of Erythema or Moist Desquamation



(Prescription)

When to Use Regenecare[®]:

- Pain due to radiation reactions
- Abrasions
- 1st and 2nd degree burns & cuts
- Dry & moist desquamation
- Reduce pain in partial and full thickness wounds
- Reduce pain from EGFR inhibitors

How to Use:

- Cleanse with **MPM Wound and Skin Cleanser** and allow drying
- Apply **Regenecare Gel** 1/4" thick & **Regenecare Gel Pad** 3Xs daily or as needed
- Apply dressings in accordance with your protocol

Benefits to Patient:

- Absorbs Minimal Drainage
- Reduces pain with 2% Lidocaine
- Reduces itching and inflammation
- Provides moisture to the skin cells and wounds
- Builds structure to damaged skin

For Treatment of Skin Necrosis or Laceration

When to Use SilverMed[™]:

- Allergic to Sulfa in Silvadine
- Skin abrasions, lacerations and tears
- Delayed-healing with pressure, foot, and leg ulcers
- Highly colonized wounds
- When an antimicrobial barrier is needed
- Help prevent Staph & Pseudomonas infection
- Help prevent MRSA & VRE
- Help prevent E. coli
- 1st & 2nd degree burns
- Regulate hydration by donating moisture

How to Use:

- Cleanse with **MPM SilverMed Antimicrobial Wound Cleanser** and allow drying
- Apply **SilverMed Hydrogel** 1/4" thick daily on the skin or throughout the wound or as needed
- Apply dressings in accordance with your protocol

Benefits to Patient:

- Does not contain Sulfa
- Reduces bioburden
- Provides long-lasting microbial barrier
- Effective against a broad spectrum of pathogenic bacteria & fungi
- Assists in prevention of contamination and infection



For Treatment of Radiation Dermatitis

When to Use CoolMagic[™] Hydrogel Dressing:

- Radiation Dermatitis
- Pressure Ulcers Stages I-II
- Partial-thickness wounds
- Draining and non-draining wounds
- 1st and 2nd degree burns
- Non-infected wounds



How to Use:

- Cleanse with **MPM Wound & Skin Cleanser** or **Antiseptic Wound Cleanser**
- Open package and remove **CoolMagic Gel Sheet**
- Remove and discard colored film backing
- Clear film backing should remain on gel sheet
- Exposed hydrogel side applied to intact skin or dermal wound
- Allow to overlap intact skin
- Hold in place with tape or stretch gauze
- Change dressing as often as necessary
- May be left on up to three days



Benefits to Patient:

- Cools live nerve endings
- Instant cooling sensation
- Minimizes damaging skin tissue upon dressing removal
- Hydrophilic – creates moist wound environment
- Absorbs three times its weight of drainage

For Treatment of Mouth Pain Oral Mucosa

When to Use OraMagic[®]Rx or OraMagicPlus[®]:

- Oral mucositis
- Aphthous ulcers
- Stomatitis
- Esophagitis
- Canker sores
- Braces
- Gum soreness
- HIV related mouth sores (**OraMagicPlus[®]**)
- Dental sensitivity (**OraMagicPlus[®]**)



(Prescription) (Non-Prescription)

How to Use:

- Step 1: Fill bottle with water to the first line and shake vigorously
- Step 2: Add water to the second fill line and shake
- Step 3: Swish one tablespoon slowly around mouth
- Step 4: Spit out remaining liquid
- Step 5: Repeat one hour after first application
- Step 6: Apply up to 4Xs daily or as directed

Benefits to Patient:

- Instant pain relief with Benzocaine (**OraMagicPlus[®]**)
- Protects and reduces further irritation and painful sensations
- Pleasant taste
- Coats injured oral mucosa
- Reduces inflammation
- Moisturizes and breaks down slough
- Relieves pain

www.mpmmed.com • (800) 232-5512

© MPM Medical, Inc. 2011. All Rights Reserved.

MPM
MEDICAL, INC.