

### BEFORE STARTING THE TEST

READ ALL THESE INSTRUCTIONS AND FAMILIARIZE YOURSELF WITH THE TEST PROCEDURE BEFORE STARTING THE TEST.

Test results will only be useful if samples are properly collected.

If you have any questions during test collection, please call your referring doctor or SIBO Clinic Canada.

DO NOT INSERT YOUR FINGER INTO THE TUBE HOLDER OF THE EASY SAMPLER BREATH
COLLECTOR AT ANY TIME; IT CONTAINS A SHARP NEEDLE THAT IS COVERED BY A GREY RUBBER
MEMBRANE, DO NOT TOUCH THIS, IT ACTS AS A SEAL TO CATCH THE BREATH EXACTLY

DO NOT LOOSEN OR REMOVE THE TOPS OF THE VACUUM-SEALED COLLECTION TUBES; THIS WILL DESTROY THE VACUUM AND MAKE THE TUBES USELESS FOR THE TEST.

IMPORTANT: ADD THE LACTULOSE TO APPROX 80Z'S OF WATER AND MIX, DRINK IT AFTER YOU TAKE THE FIRST BASE BREATH SAMPLE!

### **KIT CONTENTS**

- Easy Sampler with tube holder
- 10 vacuum sealed collection tubes
- 10 lables for the collection tubes
- 1; 15 ml dose of teva-lactulose
- 2 bubble wrap bags for the tubes after you are finished.
- Prepaid shipping information
- Instruction sheets / charting sheet / Intake form

# PLACE ONLY THE TUBES AND INTAKE FORM BACK IN THE BOX, ATTACH THE RETURN LABEL TO BOX THEN SHIP BACK.

#### TIME NEEDED FOR TESTING

- A full 12 hours on the restricted diet followed by 12 hours of fasting
- The test will take three hours to compete
- Perform the test one hour after waking
- Samples are collected every 20 minutes for 3 hours in total
- First perform the baseline breath BEFORE ingesting the substrate/lactulose then 9 breath samples AFTER ingesting the lactulose solution
- PLEASE schedule your time appropriately

### **DURING THE SAMPLE COLLECTION**

- Make sure there is a tight enough seal around the mouth piece
- 2 REGULAR breaths only, DO NOT take deep breaths, DO NOT blow hard in the breath collection device!
- 3 Once the bag is close to fully inflated push the tube in all the way, you will feel it puncture the tube hold in for 2 seconds then pull out and move on to the next tube.



### **Patient Preparation Guideline For SIBO Breath Testing**

This diet must be adhered to, do not consume anything other than whats mentioned here. Read all information below please!

### PROCEDURES, MEDICATIONS AND SUPPLEMENTS

### PRIOR TO BREATH TESTING, STOP THE FOLLOWING FOR THE REQUIRED TIMELINE:

- At least one month prior: colonoscopy, colonics, antibiotics, or antifungals stop
- PPI's are no longer required to stop, BUT if they could be stopped two days prior to testing it would be beneficial. "Consult with your practitioner"
- At least one week prior: promotility drugs or Laxatives
- One day prior: high dose magnesium or vitamin C

## ONE DAY PRIOR TO TESTING, PLEASE STOP THE FOLLOWING:

- Probiotics or prebiotics or any probiotic containing product (yogurt, Yakult, fermented foods)
- Herbal antimicrobials
- Digestive aids such as enzymes or hydrochloric acid
- Any non-essential medications and supplements including probiotics 24 prior to testing and during preparation and testing period

### Day of testing, avoid smoking and physical activity in the morning

<u>Diabetic and pregnant patients, PLEASE consult with your physician prior to prep.</u>

#### PREPARATION GUIDE

The purpose of this diet is to reduce food fermentation and therefore provide a clear response to lactulose.

Preparation begins 24 hours prior to testing in 2 stages of 12 hours. *If you suffer from constipation, please allow 48 hours prior to testing.* 

- Stage 1 consists of a restricted meal plan
- Stage 2 consists of fasting only water may be consumed

#### Note:

- Wake up at least an hour before commencing the test
- Brush teeth as per normal but rinse with an antiseptic if possible before starting the breath test
- No smoking, including second-hand smoke, for at least 1-2 hours before or at any time during testing period
- No sleeping or vigorous exercise for at least 1 hour before or at any time during testing period
- Drinking water during the breath test is allowed in moderation
- Recent antibiotic therapy or runny diarrhea may affect the results of the breath tests – \*\*please consult with your health practitioner about these conditions prior to performing the test as testing may need to be rescheduled\*\*

#### ALLOWED FOOD LIST

# Only consume the following for the first 12 hours, 48 hours if constipated, to ensure accuracy of test results:

- Baked or broiled chicken, fish or turkey (Salt and pepper for seasoning)
- Plain steamed white rice (Basmati or Jasmine)
- Hard aged cheese (Parmesan, Pecorino), unless sensitive to dairy
- Eggs
- Clear meat broth (not bone or vegetables)
- Olive oil or coconut oil 1 tablespoon only, for cooking
- NO herbal teas
- Plain water NO mineral water

#### **SAMPLE DIET**

**8AM TILL 8PM:** 12 or 48 hours of restricted meal

plan

Breakfast: Scrambled eggs

**Lunch:** Chicken with white jasmine or basmati rice **Dinner:** Fish and white jasmine or basmati rice

**Drink:** Water

7PM TILL 7AM: 12 hours fasting

\*Prior to testing, watch the instructional video on the front page of our website: www.sibocliniccanada.com



### **PATIENT INTAKE FORM**

Please read and check all the apply!

Patient Name:	Ethnicity:				
DOB:	Gender: M 🗌 F 🔲 E	Body Weight:	lbs		
Address:	City: _		_ Prov/Sate:	Postal Code/Zip:	
	Er			PRINT CLEAR	
(Optional) Clinic & I	Doctor / Practitioner Name: _				
Phone Number:	Er	nail:		does not require reference forms from practitions or doctors.	
Breath Collection D	Pate:			<u> </u>	
	?: <b>YES</b> NO lf yes, date o		Results: <b>Po</b>	sitive Negative	
Type of treatment:	Antibiotic Antimicrobia	als 🔲 Other (pl	ease specify):		
	SYMPTOM	S (Check all th	at apply)		
<ul><li>Vomiting</li><li>Diarrhea</li><li>Bloating</li><li>Constipation</li><li>Please</li></ul>	Gassy (smelling like rotten eggs) Anxiety Leaky Gut  DISEASE/DISO e list any diagnosed or susp	Fatigue	shes ntal disease & halit	_	
•					
•		•			
	Tube 1 BASE BREATH, please				
Drink Lactulose solu		Please Record Time			
Tube #2		20 min after drink			
Tube #3		40 min after drink			
Tube #4 Tube #5		60 min after drink 80 min after drink			
Tube #5		100 min after drink			
Tube #6		120 min after drink			
Tube #7		140 min after drink			
Tube #9		160 min after drink			
Tube #10		180 min after drink			
Tube #10	100 111111 411	ter drink			
	es should be received by SIB A does not diagnose or treat any				

SIBO CLINIC CANADA does not diagnose or treat any disease, aliment or other health symptoms. Any diagnosis or treatment needs to be made by your healthcare provider. The above tests are meant to aid your physician in their diagnosis, which is made in accordance with other clinical data that is unavailable to SIBO CLINIC CANADA.

SIGN:	DATE:



### **RETURN INSTRUCTIONS**

- 1 Return label is valid for 60 days.
- 2 Inside the kit are two small bubble wrap envelopes. Put the tubes in these envelopes (5 per pack) put them back in the box along wit the patient intake from. Discard the rest.
- 3 Attach the Return label to the white box (do not tape over barcode). Drop to the nearest Post Office, UPS or Fedex location, whichever label we used in your kit. Sometimes we use different carries for pricing reasons. Don't forget to keep your tracking number.

Expected arrival back to our facility, 2-5 business days. Not guaranteed with COVID. But it's still efficient.

If there are any issues, please call us directly ASAP. 506 381 9847