Drug Facts

Active ingredients (in each 20 mL)

Purpose

Acetaminophen 650mg ... Painreliever/feverreducer

Dextromethorphan HBr 20 mg......Cough suppressant

Guaifenesin 400 mg.....Expectorant

Phenylephrine HCl 10 mg.....Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough

- nasal congestion
- minor aches and pains
- sore throat
- headache
- stuffy nose
- sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersomemucusandmakecoughsmoreproductive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening
■ blisters
■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consultadoctor promptly.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor before use if you have taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

Inactive ingredients anhydrouscitric acid, cetylpyridiniumchloride, dibasic, sodium, phosphate, edetatedisodium, flavor, hydrochloric acid, polyethyleneglycol, propyleneglycol, purified water, sodiumhydroxide, sucralose

If pregnant or breast-feeding, askahealth professional before use.

Keep out of reach of children.

Overdose warning: Takingmorethan the recommended dose (overdose) maycauseliver damage. Incaseofoverdose ,getmedical help or contact a Poison Control Centerrightaway. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over:20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Inactive ingredients: ammonium, glycyrrhizate, anhydrouscitricacid, D & Cyellow no.10, edetatedisodium, FD & C blueno.1, flavors, glycerin, propyleneglycol, purifiedwater, sodium benzoate, sodiumcitrate, sorbitol, sucralose, xanthangum.

Questions? 1-866-MUCINEX (1-866-682-4639)

Other information

- each20mLcontains: sodium 12 mg
- storeat20-25°C(68-77°F)
- do notrefrigerate

