

NIGHT TIME SEVERE COLD & FLU

Drug Facts

Active ingredients Purpose (in each 20 mL)

Acetaminophen 650mg ... Painreliever/feverreducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Phenylephrine HCl 10mgNasaldecongestant
Triprolidine HCl 2.5 mg.....Antihistamine

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - itching of the nose or throat
 - sore throat
 - minor aches and pains
 - stuffy nose
 - sinus congestion and pressure
 - runny nose
 - temporarily reduces fever
 - headache
 - nasal congestion
 - sneezing
- itchy, watery eyes due to hay fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a breathing problem such as emphysema or chronic bronchitis

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product do not use more than directed

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

Drug Facts (continued)

- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20mL contains: **sodium 16 mg**
- store at 20-25°C (68-77°F) do not refrigerate

Inactive ingredients: ammonium, glycyrrhizate, anhydrous citric acid, acid ascorbic D&C yellow no. 10, edetate disodium, FD&C blue no. 1, flavors, FD&C red no. 40 glycerin, (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethylcitrate, water, xanthan gum.

Questions? 1-866-MUCINEX (1-866-682-4639)

MUCINEX®

KICKSTART SEVERE COLD & FLU

Drug Facts

Active ingredients (in each 20 mL)

Purpose

Acetaminophen 650mg ... Painreliever/feverreducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mgExpectorant
Phenylephrine HCl 10mgNasaldecongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - itching of the nose or throat
 - sore throat
 - minor aches and pains
 - stuffy nose
 - sinus congestion and pressure
 - runny nose
 - temporarily reduces fever
 - headache
 - nasal congestion
 - sneezing
- helps loosen-phlegm(mucus) and thin bronchial secretions to rid the bronchial passage ways of bother some mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe,persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, orvomiting, consult a doctor promptly.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease),or for 2weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product ■ do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,
askahealth professional before use.

Drug Facts (continued)

Keep out of reach of children.

Overdose warning: Takingmorethan the recommended dose (overdose) maycauseliver damage. Incaseofoverdose ,getmedical help or contact a Poison Control Centerrightaway. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 12 mg**
- storeat 20-25°C(68-77°F) ■ do not refrigerate

Inactive ingredients : ammonium, glycyrrhizate, anhydrous citric acid, acid ascorbic D&C yellow no.10, edetate disodium, FD&C blue no. 1, flavors, glycerin, propylene glycol, purified water, sodium benzoate, citrate,sorbitol, sucralose, xanthan gum.

Questions? 1-866-MUCINEX (1-866-682-4639)

***FAST-MAXKICKSTARTSEVERECOLD& FLU**
helps to relieve these symptoms day or nigh zt; FAST- MAX NIGHT TIME SEVERE COLD & FLU helps to relieve.

Do not take Mucinex Fast -Max Kick start Severe Cold & Flu and MucinexFast-MaxNightTimeSevere Cold & Flu at the same time. Always wait at least 4 hours before taking another dose of Mucinexliquid.

MUCINEX®